


Advanced Functions

Course Notes

Chapter 3 – Polynomial Equations and Inequalities

We will learn

- *how to find solutions to polynomial equations using tech and using algebraic techniques*
- *how to solve polynomial inequalities with and without tech*
- *how to apply the techniques and concepts to solve problems involving polynomial models*



Chapter 3 – Polynomial Equations and Inequalities

Contents with suggested problems from the Nelson Textbook (Chapter 4). You are welcome to ask for help, from myself or your peers, with any of the following problems. They will be handed in on the day of the Unit Test as a homework check.

3.1 Solving Polynomial Equations – Pg 57 - 61

Pg. 204 – 206 #1, 2, 6 – 8, 10 – 12, 14, 15

3.2 Linear Inequalities – Pg 63 – 66

Pg. 213 – 215 #1, 2, 4, 5, 7, 9, 13

3.3 Solving Polynomial Inequalities – Pg 67 – 70

Pg. 225 – 228 #2, 5 – 7, 10 – 13

3.1 Solving Polynomial Equations

Learning Goal: We are learning to solve polynomial equations using a variety of strategies.

Before embarking on this wonderful journey, it seems to me that it would be prudent to make some (seemingly silly) opening statements.

Seemingly Silly Opening Statements

- 1) Polynomial **equations ARE NOT** polynomial **functions!**
- 2) Solving any equation **MEANS** finding a **SOLUTION** (if a solution exists)!
- 3) Solving a polynomial equation is **ALWAYS** equivalent to finding the **zeros** of some polynomial function!

Example 3.1.1 (back to Grade 9)

Solve the linear equation

$$3(x-5)+2=5x+6$$

$$3x-15+2=5x+6$$

$$3x-13=5x+6$$

$$-2x=19$$

$$x = \frac{-19}{2}$$

Example 3.1.2 (remember grade 11?)

Solve the quadratic equation

$$5x(x-1)+7=2x^2+9$$

$$5x^2 - 5x + 7 = 2x^2 + 9$$

$$3x^2 - 5x - 2 = 0$$

$$\underline{3x^2 - 6x} + \underline{1x - 2} = 0$$

$$(3x+1)(x-2) = 0$$

Goal:

$$\text{"stuff"} = 0$$

- ① Quadratic Formula
- ② Factoring!!

$$\therefore x = -\frac{1}{3} \text{ and } x = 2$$

Geometrically speaking, solving a quadratic **equation** is equivalent to finding the zeros of a quadratic **function**.

Solving the **equation** in **Example 3.1.2** means the same thing as finding the zeros of the **function**

Note further that quadratic **functions** can have



Thus quadratic **equations** can have **2 solutions**, **1 solution** or **no solutions**!

Comments about Higher Order Polynomial Equations

Consider the cubic **EQUATION** $x^3 + 2x^2 - 5x + 1 = 0$.

Q. How many zeros can this equation have?

Ans.

1, 2, or 3

Odd's must have at least one solution.

Consider the quartic equation $4x^4 - 3x^3 + 5x^2 - 3x + 1 = 0$.

Q. How many zeros can this equation have?

Ans.

0, 1, 2, 3, or 4.

Even's could have no solutions.

Example 3.1.3

Solve the polynomial equation by factoring:

$$4x^3 - 3x = 1$$

$$4x^3 - 3x - 1 = 0 \quad \text{try } \pm 1$$

$$\text{Try } f(1) = 4(1)^3 - 3(1) - 1$$

$$= 0 \quad \therefore x-1 \text{ is a factor.}$$

$$\begin{array}{r|rrrrr} 1 & 4 & 0 & -3 & -1 & \\ & & 4 & 4 & 1 & \\ \hline & 4 & 4 & 1 & 0 & \\ & x^2 & x^1 & x^0 & & \end{array}$$

$$\begin{aligned} \therefore 4x^3 - 3x - 1 &= (x-1)(4x^2 + 4x + 1) \\ &= (x-1)(2x+1)^2 \end{aligned}$$

$$\therefore x = 1 \text{ and } x = -\frac{1}{2}$$

Example 3.1.4

Solve the equation by factoring:

$$12x^4 + 16x^3 - 11x = 13x^2 - 6$$

$$12x^4 + 16x^3 - 13x^2 - 11x + 6 = 0$$

$$\text{Try } f(-1) = 12(-1)^4 + 16(-1)^3 - 13(-1)^2 - 11(-1) + 6$$

$$= 0! \quad \therefore (x+1) \text{ is a factor}$$

$$\begin{array}{r|rrrrr} -1 & 12 & 16 & -13 & -11 & 6 \\ & & -12 & -4 & 17 & -6 \\ \hline & 12 & 4 & -17 & 6 & 0 \end{array}$$

$$\therefore (x+1)(12x^3 + 4x^2 - 17x + 6)$$

$$\text{Let } g(x) = 12x^3 + 4x^2 - 17x + 6$$

Test $\pm 1, \pm 2, \pm 3, \pm 6$

$$g(\pm 1) \neq 0, g(\pm 2) \neq 0, g(\pm 3) \neq 0, g(\pm 6) \neq 0$$

Note: Solving Polynomial Equations requires writing the equation in **Standard Form**, which is: "polynomial = 0"

Rational Zero Test

Consider $12x^3 + 4x^2 - 17x + 6 = 0$.

We now, when using the factor theorem, will "test for zeros" using 2 steps:

- 1) Test for integer zeros using factors of the constant term.

$$\pm 1, \pm 2, \pm 3, \pm 6$$

- 2) Test for rational zeros, where we consider $x = \frac{b}{a}$

$$\pm \frac{1}{2}, \pm \frac{1}{3}, \pm \frac{1}{4}, \pm \frac{1}{6}, \pm \frac{1}{12}, \pm \frac{2}{3}, \pm \frac{3}{2}, \pm \frac{3}{4}$$

The possible rational zeros are:

24 possible factors.

Back to Example 3.1.4

For $g(x) = 12x^3 + 4x^2 - 17x + 6$ the possible rational zeros are:

$$\text{Test } g\left(\frac{1}{2}\right) = 12\left(\frac{1}{2}\right)^3 + 4\left(\frac{1}{2}\right)^2 - 17\left(\frac{1}{2}\right) + 6$$

$$= 12\left(\frac{1}{8}\right) + 4\left(\frac{1}{4}\right) - \frac{17}{2} + 6$$

$$= \frac{3}{2} + 1 - \frac{17}{2} + 6 = \frac{3}{2} + \frac{2}{2} - \frac{17}{2} + \frac{12}{2} = \frac{0}{2} = 0!$$

$$\begin{array}{r|rrrr} \frac{1}{2} & 12 & 4 & -17 & 6 \\ & & 6 & 5 & -6 \\ \hline & 12 & 10 & -12 & 0 \end{array}$$

$\rightarrow \div 2$

$$6 \quad 5 \quad -6$$

$$\begin{aligned} \therefore &= (2x-1)(6x^2+5x-6) \\ &= (2x-1)\left(\frac{6x^2-4x+9x-6}{2 \quad +3}\right) \\ &= (2x-1)(2x+3)(3x-2) \end{aligned}$$

$$\begin{array}{l} \textcircled{x} -36 \\ \textcircled{+} +5 \\ 9,4 \end{array}$$

$$\therefore x = \frac{1}{2}, x = -\frac{3}{2}, x = \frac{2}{3}$$

Example 3.1.5Solve the equation $3x^3 - 4x + 2 = 0$. $\pm 1, \pm 2$ $\pm 1, \pm 2$ $\pm \frac{1}{3}, \pm \frac{2}{3}$

$$\begin{aligned} \text{Test: } f\left(\frac{1}{3}\right) &= 3\left(\frac{1}{3}\right)^3 - 4\left(\frac{1}{3}\right) + 2 \\ &= \frac{3}{27} - \frac{4}{3} + 2 \neq 0 \end{aligned}$$

As it turns out... $f(\pm 1) \neq 0$, $f(\pm 2) \neq 0$, $f(\pm \frac{1}{3}) \neq 0$
 $f(\pm \frac{2}{3}) \neq 0$.

$\therefore f(x)$ does not factor.

However, this is a cubic fn, and therefore it must
 have at least one.

Use technology,...

$x = -1.352$ by graphing...

Success Criteria:

- I can solve polynomial equations algebraically (by factoring) AND graphically
- I can recognize that only SOME polynomial equations can be solved by factoring
- I can recognize that some solutions may not make sense in the context of the question

3.2 Linear Inequalities

Learning Goal: We are learning to solve linear inequalities.

Once again, it seems a good idea to begin with a couple of opening statements.

Absolutely Non-Silly Opening Statements

- 1) The **algebra** of inequalities is the **SAME** as the algebra on equality (i.e. solving equations), with two exceptions:

- a) If you multiply or divide by a negative, then you must flip the inequality sign.

- b) We can have 2 sided inequalities – e.g.

$$3 \leq x \leq 5 \quad \text{or} \quad -3 > x > 8$$

- 2) The **Solution Set** of inequalities is infinite.

$$\begin{array}{l} -2x \leq 4 \\ \hline x \geq -2 \end{array}$$

$$\begin{array}{l} +2x \quad -4 \quad +4 \quad +2x \\ -2x < 4 \\ -4 < 2x \\ \hline -2 < x \end{array}$$

Example 3.2.1

Solve the (linear) inequality $3x - 2 > 4$.

① Set Notation:

$$\{x \in \mathbb{R} \mid x > 2\}$$


$$\frac{3x}{3} > \frac{6}{3}$$

$$x > 2$$

② Interval Notation

$$x \in (2, \infty)$$

③ Graphing on a # line.



Do the "math" everywhere

Example 3.2.2

Solve the two sided inequality $-2 > -4x + 5 \geq -3$.


$$\frac{-7}{-4} > \frac{-4x}{-4} \geq \frac{-8}{-4}$$

$$\frac{7}{4} < x \leq 2$$

① Interval:

$$x \in \left(\frac{7}{4}, 2\right]$$

② Graph



Example 3.2.3

Solve $5 \leq 3(x-2) - 4(x+3) \leq 12$

$$5 \leq 3x - 6 - 4x - 12 \leq 12$$

$$\overset{+18}{5} \leq -x - 18 \leq \overset{+18}{12}$$

$$\frac{23}{-1} \leq \frac{-x}{-1} \leq \frac{30}{-1}$$

$$-23 \geq x \geq -30$$

arrange small to big...

$$-30 \leq x \leq -23$$

$$x \in [-30, -23]$$

Example 3.2.4

Write the following sketch of a solution set in interval and set notation:




Figure 3.2.4

$$\text{Interval: } x \in (-\infty, -4) \cup [3, \infty)$$

$$\text{Set: } \{x \in \mathbb{R} \mid -4 > x \geq 3\}$$

Graphical Views of (non-linear) Polynomial Inequalities

(the Algebra is tough...)

Example 3.2.5

Consider the sketch of the graph of some mystery cubic function.

Q. When (or better WHERE) are the *functional values* positive?

↖ ↗




Figure 3.2.5

Example 3.2.6

Consider the sketch of the quartic $g(x)$, and determine where

- a) $g(x) \leq 0$
- b) $g(x) > 2$
- c) $-1 \leq g(x) \leq 2$

a) $x \in [-4, 1] \cup [3]$
 $\{3\}$

b) $x \in (-\infty, -5) \cup (5, \infty)$

c) $x \in [-5, -3] \cup [0, 5]$




Figure 3.2.6

Success Criteria:

- I can solve a linear inequality by using inverse operations
- I can recognize that when you multiply/divide by a negative number, you MUST reverse the inequality sign
- I can recognize that linear inequalities have many solutions
- I can express the solution to a linear inequality on a number line

3.3 Solving Polynomial Inequalities

Learning Goal: We are learning to solve polynomial inequalities.

For this section, no opening statements are required....

Non-Required Opening Statement

Solving non-linear polynomial inequalities can be accomplished in two ways:

- 1) Graphically (sometimes called Geometrically)
- 2) Algebraically (which tends to be more useful)


Example 3.3.1

Solve $(2x-1)(x-2)(x+3) \geq 0$

REMEMBER: *FACTORED FORM IS YOUR FRIEND*

Graphically:

$$y = (2x-1)(x-2)(x+3) = 6$$



Note: Solving an inequality graphically is

rather easy, **BUT** solving algebraically
may require more work.

$$x \in [-3, \frac{1}{2}] \cup [2, \infty)$$

Example 3.3.1 (Continued)

$$\text{Solve } (2x-1)(x-2)(x+3) \geq 0$$

Algebraically

For this technique we will construct an “**Interval Chart**”, which can also be thought of as a “**table of signs**” (and wonders?)

Note: It is often helpful to remember that in mathematics we are dealing with **NUMBERS**.

Numbers have signs: **Positive** or **Negative**

e.g. $(x-2)$ is a **NUMBER** whose sign switches from +’ve to -’ve at $x=2$ (i.e. the sign switches at the zero of the factor)

The Interval Chart looks like:

Intervals	Split the Domain $(-\infty, \infty)$ at all ZEROS of the Factors		
Test Values	Choose a Domain	value inside each	Interval
Sign on 1 st Factor			
Sign on 2 nd Factor			
Sign on 3 rd Factor			
Sign on the Product of Factors	Find the Intervals	with the sign we	want to answer the question

For our problem above, our chart will look like:

$$x = -3, \frac{1}{2}, 2$$

Intervals	$(-\infty, -3)$	$(-3, \frac{1}{2})$	$(\frac{1}{2}, 2)$	$(2, \infty)$
Test Values	-4	0	1	3 or 3,000,000
$x+3$	—	+	+	+
$2x-1$	—	—	+	+
$x-2$	—	—	—	+
Product	—	+	—	+

$$\therefore x \in [-3, \frac{1}{2}] \cup [2, \infty)$$

$\pm 1, \pm 2, \pm 3, \pm 6, \pm 9, \pm 18$

Example 3.3.2

Solve algebraically $4x^4 + 16x^3 + x^2 - 39x - 18 < 0$.

Let $f(x) = 4x^4 + 16x^3 + x^2 - 39x - 18$

Wait a second....where is your friend and mine...

Factored Form!!

$$f(-2) = 4(-2)^4 + 16(-2)^3 + (-2)^2 - 39(-2) - 18$$

$$= 64 - 128 + 4 + 78 - 18$$

$f(-2) = 0!!$ Thanks Jake $\therefore (x+2)$ is a factor.

$$\begin{array}{r|rrrrr} -2 & 4 & 16 & 1 & -39 & -18 \\ & \downarrow & -8 & -16 & 30 & 18 \\ \hline & 4 & 8 & -15 & -9 & 0 \end{array}$$

$\therefore (x+2)(4x^3 + 8x^2 - 15x - 9) < 0$

$\pm 1, \pm 3, \pm 9$

Let $g(x) = 4x^3 + 8x^2 - 15x - 9$

$$g(-3) = 4(-3)^3 + 8(-3)^2 - 15(-3) - 9$$

$$= -108 + 72 + 45 - 9$$

$= 0!!$ Thanks Alex.

$\therefore (x+3)$ is a factor.

$$\begin{array}{r|rrrr} -3 & 4 & 8 & -15 & -9 \\ & \downarrow & -12 & 12 & 9 \\ \hline & 4 & -4 & -3 & 0 \end{array}$$

$\therefore (x+2)(x+3)(4x^2 - 4x - 3) < 0$

$= (x+2)(x+3)(4x^2 - 6x + 2x - 3) < 0$

$m = -12$
 $A = -4$
 $-6, 2$

$= (x+2)(x+3)(2x+1)(2x-3) < 0$

$x = -3, -2, -\frac{1}{2}, \frac{3}{2}$

Intervals	$(-\infty, -3)$	$(-3, -2)$	$(-2, -\frac{1}{2})$	$(-\frac{1}{2}, \frac{3}{2})$	$(\frac{3}{2}, \infty)$
T.V.	-4	-2.5	-1	0	2
$x+3$	-	+	+	+	+
$x+2$	-	-	+	+	+
$2x+1$	-	-	-	+	+
$2x-3$	-	-	-	-	+
Product	+	-	+	-	+

$$\therefore 4x^4 + 16x^3 + x^2 - 39x - 18 < 0$$

$$\text{when } x \in (-3, -2) \cup (-\frac{1}{2}, \frac{3}{2})$$

Success Criteria:

- I can solve polynomial inequalities algebraically by
 1. Moving all terms to one side of the inequality
 2. Factoring to find the zeros of the corresponding polynomial
 3. Creating a number line, graph, or an interval chart
 4. Determining the intervals on which the polynomial is positive or negative
- I can solve polynomial inequalities graphically