


# Advanced Functions

## Course Notes

## Chapter 1 – Functions

**Learning Goals:** We are learning to...

- identify the characteristics of functions and how to represent them
- apply transformations to parent functions and how to use transformations for sketching
- to determine the inverse of a function



# Chapter 1 – Introduction to Functions

*Contents with suggested problems from the Nelson Textbook. These problems are not going to be checked, but you can ask me any questions about them that you like.*

## **1.1 Functions – Pg 1 - 3**

Read Example 3 on Page 9 - Pg. 11 – 13 #1 – 3, 5, 6, 7b-f, 9, 10, 12

## **1.2 Properties of Functions – Pg 4 – 12**

Pg. 23 – 24 #5, 7 – 11 (*1.3 in Nelson Text*)

## **1.3 Transformations of Functions Review – Pg 13 - 15**

Worksheet and graphs

## **1.4 Inverses of Functions – Pg 16 - 20**

Pg. 43 – 45 #2 – 4, 7, 9, 12, 13, 15 (*1.5 in Nelson*)

## **1.5 Piecewise Defined Functions – Pg 21 - 25**

(Abs. Value.) Pg. 16 #2, 4 – 8 (think about transformations!), 10 (*1.2*)

(Piecewise) Pg. 51 – 53 #1 – 5, 7 – 9 (*1.6*)

# 1.1 Functions

**Learning Goal:** We are learning to represent and describe functions and their characteristics.

There are some people who argue that mathematics has just two basic building blocks: Numbers and Operations. This course is concerned with functions which can be considered number generators. A function takes a given number, and using mathematical operations generates another number. We will be examining the **relationship** between the given numbers, and the generated numbers for various functions.



## Definition 1.1.1

A **Function** is an algebraic rule which assigns exactly one element in a set called the range to each element in a set called the domain.  $\rightarrow$   $x$ -values  $y$ -values

$\rightarrow$  each  $x$ -value produces only one  $y$ -value.


## Pictures

### Vertical Line Test




This is a function as it passes the VLT


### Arrow Diagrams Mapping diagrams



2 Is a function



Not a function because 1 produced 8 and 10



Is a function

$$f(x) = \sqrt{x-1}$$

### Definition 1.1.2

**Domain** of a Function: a set of "x" values which makes sense when plugged into the function

**Range** of a Function

the set of functional values (y) which are produced from the domain values

### Function Notation

We use the notation  $f(x)$  to "name" a function. This notation is powerful because it contains both the domain and the range. For example we might write  $f(2)$ , which shows that the domain value is  $x=2$ , and that the range value (which we must calculate) is denoted  $f(2)=8$

Domain  $\rightarrow$  Range  $(2, 8)$   $f(x) = y$

### Definition 1.1.3

The **Graph** of a function is a set of points  $(x, f(x))$ , denoted

$$as \ f(x) = \{ \underbrace{(x, f(x))}_{\text{points such that}} \mid x \in D_f \}$$

belongs to.

### Example 1.1.1

Given the graph of the function  $f(x) = \{(3, 4), (2, -1), (7, 8), (4, 2), (5, 4)\}$  determine:

- $D_f = \{3, 2, 7, 4, 5\}$
- $R_f = \{4, -1, 8, 2\}$
- Is  $f(x)$  a function? YES. Each x-value is listed once

### Example 1.1.2

Consider the sketch of the graph of  $g(x)$ , and determine:

- $D_g = \{x \in \mathbb{R} \mid -4 \leq x \leq 5\}$
- $R_g = \{y \in \mathbb{R} \mid -2 \leq y \leq 3\}$
- Is  $g(x)$  a function?

Not a function. Failed the VLT




Figure 1.1.2

Note: In the above examples we have seen functions (and non-functions which we call relations) depicted graphically and numerically. We now turn to algebraic representations of functions. It is much more difficult to determine domain and range for functions given in an algebraic form, but the algebraic form is incredibly useful!


### Example 1.1.3

State the domain and range of the functions given in algebraic form.

a)  $f(x) = 3\cos(2x)$

b)  $g(t) = (t-2)^2 + 1$


c)  $h(x) = \frac{2}{x-1}$



$$D_f = \{x \in \mathbb{R}\}$$


$$\rightarrow R_f = \{f(x) \in \mathbb{R} \mid -3 \leq f(x) \leq 3\}$$

$$= f(x) \in [-3, 3]$$



$$D_g = \{t \in \mathbb{R}\}$$

$$R_g = \{g(t) \in \mathbb{R} \mid g(t) \geq 1\}$$



$$D_h = \{x \in \mathbb{R} \mid x \neq 1\}$$

$$R_h = \{h(x) \in \mathbb{R} \mid h(x) \neq 0\}$$

### Notations for Domain and Range

Assumes Real numbers

Interval Notation

$$x \in [-4, 5]$$

Set Notation

$$\text{uses } \{ \}$$

~~Pseudo set Notation~~

### Success Criteria

- I can use function notation to represent the values of a function
- I can apply the vertical line test
- I can identify the domain and range for different types of functions
- I can recognize and apply restrictions on the domains of functions

We use square brackets and round brackets.

[ ]

when the end point is included

• or  $\leq, \geq$

( )

when the end point is not included

o or  $<, >$

## 1.2 Properties of Functions

**Learning Goal:** We are learning to compare and contrast the properties and characteristics of various types of functions.


Recall that we define the graph of a function to be the SET of Ordered Pairs:

We can visualize the graph of a function by plotting its ordered pairs on the Cartesian axes.

### Example 1.2.1

e.g.  $f(x) = x^2$  has the graph

and looks like



## Characteristics of a Function's Graph

Over the course we will be studying Polynomial, Rational, Trigonometric, Exponential and Logarithmic Functions. For now we are focussed on Polynomial and Rational Functions, but for each type of function we will try and understand various functional (fnal) behaviours (or characteristics).

The characteristics (behaviours) we are primarily interested in studying are:

1. Domain and Range
2. Axis Intercepts ( $x$  and  $y$  intercepts)
3. Intervals of Increase/Decrease
4. Continuity vs. Discontinuity
5. Function end behaviors

Note: Generally a geometric point of view will just mean that we'll look at pictures, but Geometry is actually ***much*** deeper than that!

## Intervals of Increase and Decrease

We will examine (when possible) functional behaviour from both algebraic and geometric points of view.


### Definition 1.2.1

A function  $f(x)$  is said to be increasing on the open interval  $(a, b)$  when

$$f(a) < f(b)$$

A function  $f(x)$  is said to be decreasing on the open interval  $(a, b)$  when

$$f(a) > f(b)$$



Note the difference between open and closed intervals:


An open interval does not include endpoints  $(a,b)$

A closed interval does include endpoints  $[a,b]$

### Example 1.2.2

Consider the function  $f(x)$ , represented graphically:

Determine where  $f(x)$  is increasing and decreasing.



Increasing on:  $(-\infty, 0) \cup (2, \infty)$

Decreasing on:  $(0, 2)$





# Continuity

For the time being we will consider a (quite) rough definition of what it means for a function to be continuous. In fact, we will see that understanding what it means for a function to be discontinuous may be more helpful for now. In the course *Calculus and Vectors*, a formal, algebraic definition of continuity will be considered.

## Rough Definition

A function  $f(x)$  is **continuous** (cts) on its domain  $D_f$  if *when sketching, you do not lift your pen/pencil.*


Pictures



There are 3 types of **discontinuities**:

1)

*Jump*




*Jump at  $x=5$*

2)


Hole

$$f(x) = \frac{x^2 - 16}{x - 4}$$

$$= \frac{(x-4)(x+4)}{x-4}$$



3)



Infinite (Asymptotic)

## End Behaviour of Functions

Here we are concerned with how the function is behaving as  $x$  gets

**HUGE**

As  $x$  gets **HUGE** (which we write  $x \rightarrow \infty$ , or  $x \rightarrow -\infty$ )

**HUGE**

↑  
approaches

$f(x)$  or  $y$ 's

the **functional values** (for whatever function we are studying) can do one of three things:

1) goes to infinity





e.g. as  $x \rightarrow \pm\infty$ ,  $f(x) \rightarrow \infty$   
or  
 $-\infty$

2) Settles on one specific value.

3) Oscillate, bounce between values

Pictures:

Black Box of Mystery



### Success Criteria

- I can identify types of functions based on their graphical characteristics
- I can use different characteristics (intervals of increase/decrease, ~~odd/even~~, end behaviour, continuity) to help me identify types of functions

# 1.3 Transformations of Functions

**Learning Goal:** We are learning to apply transformations to parent functions and how to use transformations for sketching

This section is pure review of material from Grade 11. If you've forgotten certain aspects of the concepts, ask for help. Recall that there are three basic transformations of functions. You've probably heard of Flips, Stretches and Shifts. More formal mathematical terms would be Reflections, Dilations and Translations, respectively. Recall also that transformations can occur both vertically and horizontally.

## Definition 1.3.1

Given a function  $f(x)$ , then we denote transformations to  $f(x)$  as

$$F(x) = a f(k(x-d)) + c$$

$\uparrow$   $\uparrow$   
 $\text{factored out}$

$a$  = vertical stretch

If  $a < 0$ , there is a flip along the  $x$ -axis

→ multiply the  $y$ 's by " $a$ "

$k$  = horizontal stretch

If  $k < 0$ , there is a flip along the  $y$ -axis

→ multiply the  $x$ 's by " $\frac{1}{k}$ "

$d$  = horizontal shift

→ add or subtract the stretched  $x$ 's by " $d$ "

$c$  = vertical shift

→ add or subtract the stretched  $y$ 's by " $c$ "

## Success Criteria:

- I can use the value of  $a$  to determine if there is a vertical stretch/reflection in the  $x$ -axis
- I can use the value of  $k$  to determine if there is a horizontal stretch/reflection in the  $y$ -axis
- I can use the value of  $d$  to determine if there is a horizontal translation
- I can use the value of  $c$  to determine if there is a vertical translation

## 1.4 Inverses of Functions

**Learning Goal:** We are learning to determine the equation of an inverse relation and the conditions for an inverse relation to be a function.

The inestimable William Groot has a saying:

*An Inverse Relation is an UNDO*


### Definition 1.4.1

A **relation** is simply an algebraic relationship between domain values and range values.

Note: All functions are relations, but not all relations are functions

e.g.  $x^2 + y^2 = 25$  is a relation, but it is not a function (it's a circle and so doesn't pass the VLT)

Consider the Arrow Diagram:



$\sin(x)$   
 $\sin^{-1}(x)$

### Big Concept

To determine the inverse of a function,  
just swap  $x$  and  $y$

### Example 1.4.1

Given the graph of  $f(x)$  determine:  $D_f, R_f, f^{-1}(x), D_{f^{-1}}, R_{f^{-1}}$

$$f(x) = \{(2, 3), (4, 2), (5, 6), (6, 2)\}$$

$$D_f = \{2, 4, 5, 6\} = R_{f^{-1}}$$


$$R_f = \{3, 2, 6\} = D_{f^{-1}}$$

$$f^{-1}(x) = \{(3, 2), (2, 4), (6, 5), (2, 6)\}$$


$f^{-1}(x)$  is not a function.

### Horizontal Line Test

Consider the Sketches



Fails the HLT, meaning that the inverse is not a function



Passes the HLT, meaning that the inverse is a function




## Determining the Inverse of a Function


We can determine the inverse of some given function in either of two ways: Graphically and Algebraically.

### Function Inverses Graphically

$$f(x) = x^2$$



Note: Finding a function inverse graphically is not a very useful method, but it can be instructive.



→ Flip the coordinates of every point

→ Inverses are always symmetric along  $y=x$

Is there a way to force  $f^{-1}(x)$  to be a function?

YES! Restricting the Domain on  $f(x)$

- do the inverse on only part of the domain
- restrict up to and including the turning point.

For  $f(x) = x^2$ , use either  $(-\infty, 0]$  or  $[0, \infty)$

## Function Inverses Algebraically

Determining algebraic representations of inverse relations for given functions can be done in (at least) two ways:

- 1) Use algebra in a "brute force" manner (keeping in mind the Big Concept)
- 2) Use ~~transformations~~ (keeping in mind "inverse operations")  
*Hagen's Method.*

### Example 1.4.2

Determine the inverse of  $f(x) = 2\sqrt{\frac{1}{3}(x-1)} + 2$ . State the domain and range of both the function and its inverse.

Here we will use "brute force".  
Method:

- 1) Switch  $x$  and  $f(x)$ , and  
~~call  $f(x)$   $y$~~
- 2) Solve for  ~~$f(x)$~~   $y$
- 3) turn  $y$  into  $f^{-1}(x)$

$$x = 2\sqrt{\frac{1}{3}(y-1)} + 2$$

$$x - 2 = 2\sqrt{\frac{1}{3}(y-1)}$$

$$\frac{x-2}{2} = \sqrt{\frac{1}{3}(y-1)}$$

$$3\left(\frac{x-2}{2}\right)^2 = \cancel{3}\frac{1}{3}(y-1)$$

$$3\left(\frac{x-2}{2}\right)^2 = y-1$$

$$3\left(\frac{x-2}{2}\right)^2 + 1 = y$$

$$\therefore f^{-1}(x) = 3\left(\frac{x-2}{2}\right)^2 + 1$$

### Example 1.4.3

Using ~~transformations~~ *Hagen's method!* determine the inverse of  $f(x) = 2\sqrt{\frac{1}{3}(x-1)} + 2$ .

Pretend you plug a # into  $x$ . How would you evaluate it?

$$\begin{array}{l|l} -1 & +1 \\ x \frac{1}{3} & \div \frac{1}{3} \text{ or } \times 3 \\ \sqrt{\phantom{x}} & (\phantom{x})^2 \\ \times 2 & \div 2 \\ +2 & -2 \end{array}$$

$$f^{-1}(x) = 3\left(\frac{x-2}{2}\right)^2 + 1$$



### Example 1.4.4


a) Determine the inverse of  $g(x) = -2(x-1)^2 + 3$ .

b) Note that the natural domain of  $g(x)$  is  $(-\infty, \infty)$ . However,  $g(x)$  does not pass the HLT so its inverse is not a function. Determine a restricted domain for  $g(x)$  so that  $g^{-1}(x)$  is a function.

a)

$$\begin{array}{l} -1 \\ (-)^2 \\ x - 2 \\ + 3 \end{array} \quad \begin{array}{l} + 1 \\ \div -2 \\ - 3 \end{array}$$

$$g^{-1}(x) = \sqrt{\frac{x-3}{-2}} + 1$$



### Example 1.4.5

Given  $f(x) = kx^2 - 3$  and given  $f^{-1}(5) = 2$ , find  $k$ .

Try ~~different~~ methods:

$$\begin{cases} 5 = k(2)^2 - 3 \\ 8 = k(4) \\ 2 = k \end{cases}$$

$$\begin{aligned} f^{-1}(x) &\Rightarrow (5, 2) \\ f(x) &= (2, 5) \\ &\quad \times \quad f(k) \end{aligned}$$

$$\therefore f(x) = 2x^2 - 3$$

### Success Criteria:

- I can determine the equation of an inverse function using various methods
- I can determine whether an inverse relation is a function, and whether or not the domain needs to be restricted

# 1.5 Piecewise Defined Functions

**Learning Goals:** We are learning to understand, interpret, and graph situations that are described by piecewise functions; and learning the properties of the absolute value function.

Some aspects of “reality” exhibit different (as opposed to changing)

behaviors

To capture those different behaviors mathematically may require using different

functions

over different

pieces / intervals

of the domain.

## Absolute Value

Before discussing piecewise defined functions in general, we will first review the concept of *absolute value*.

### Definition 1.5.1

The absolute *value* of a number,  $x$ , is given by


$$|x| = \begin{cases} x, & x \geq 0 \quad [0, \infty) \\ -x, & x < 0 \quad (-\infty, 0) \end{cases}$$

e.g.'s

$$|22| = 22$$

$$|-8| = -(-8) = 8$$

$$|8-13| = |-5| = 5$$




## Absolute Value Functions

We can define the function which returns the absolute value for any given number as

$$f(x) = |x| = \begin{cases} x, & [0, \infty) \\ -x, & (-\infty, 0) \end{cases}$$

Picture



(Two behaviours!)


We can go further and define functions which return the absolute value for more complicated expressions.

e.g. Sketch  $g(x) = |x^2 - 1|$  (note:  $g(x)$  takes the absolute value of the **functional values** for the “basic” function  $f(x) = x^2 - 1$ )

→ Draw  $f(x) = x^2 - 1$

→ Next, take the absolute value of the  $f(x)$  values.

$$g(x) = \begin{cases} x^2 - 1, & (-\infty, -1] \\ -(x^2 - 1), & (-1, 1) \\ x^2 - 1, & [1, \infty) \end{cases}$$



(Three functional behaviours)

## Absolute Value and Domain Intervals (and Quadratic Inequalities)

e.g.'s Sketch the solution sets of the following inequalities:

a)  $x > -1$

b)  $x \leq 2$

c)  $1 < x \leq 4$

d)  $-2 < x < 2$

Note the symmetry in part d)! Sometimes it's useful to think of absolute value as

Using the above notion we can thus use absolute value to denote the interval  $-2 < x < 2$  as



e.g. Solve the quadratic equation

$$x^2 = 4$$

e.g. Solve the quadratic inequalities, and sketch the solution sets:

a)  $x^2 < 4$

b)  $x^2 \geq 3$


And now we return our attention to general **Piecewise Defined Functions**

### Example 1.5.1

You are saving for university, and place \$1000 into a sock every six months. After 18 months you wake up and put the money in your sock into an interest bearing bank account. You continue making deposits. Give a graphical representation of this situation.

*What is the behaviour of the amount of money you have saved? How is the behaviour changing?*

Note the notation we use for piecewise defined functions. Each functional behaviour has a mathematical representation, defined over its own piece of the domain (just like the Absolute Value function we considered earlier).



**Example 1.5.2**

Determine the graphical representation for:


$$f(x) = \begin{cases} x^2, & x \in [-1, 2) \\ 2, & x \in [2, 3] \\ x+1, & x \in (3, \infty) \end{cases}$$

$$(-1)^2 = 1 \quad (2)^2 = 4$$

$$3+1=4$$

$$4+1=5$$

Continues to infinity



**Example 1.5.3**

Determine a possible algebraic representation which describes the given functional behaviour.




Figure 1.5.3

**Success Criteria:**

- I can absolutely understand the absolute value function
- I can graph the piecewise function by graphing each piece over the given interval
- I can determine if a piecewise function is continuous or not