

## Math 9 – Unit 1: Real Numbers

Name: Mr. Hagen

### Lesson #1: Rational and Irrational Numbers

Date: Sept. 11, 2017

Welcome to the wonderful and beautiful world of Mathematics. Math is a language with its own syntax, grammar, and rules. Also, for Math to be readable and elegant (yes, it can be elegant), it needs to be written in a certain way. It is essential that you learn and adapt to this structure. First, we begin by looking at real numbers.

A **real number** is *any number that you can think of.*

ex:  $5, -7, 0.08125, \pi = 3.14159...$

Another set of numbers of interests are **integers**. Integers are *non decimal numbers*

ex:  $5, -7$   $\rightarrow$  "whole number"

Within the real numbers are two different sets (or types) of numbers:

A **rational number** is: *a number which can be written as the ratio of two integers, a and b, but b cannot equal zero.*

$$\rightarrow \frac{a}{b}$$

*A rational number is a number that can be written as a fraction.*

An **irrational number** is:

*a number that cannot be written as a fraction.*

ex:  $\pi = 3.14159.....$

State if the following are rational or irrational:

a) $\frac{1}{2}$ rational	b) $\frac{-3}{0}$ neither $\rightarrow$ not a #	c) $\frac{-0}{4}$ rational	d) $\sqrt{5}$ irrational	e) $\sqrt{-9}$ You cannot square root negatives $3 \times 3 = 9$ $-3 \times -3 = 9$
------------------------------	---	-------------------------------	-----------------------------	--

Rational numbers can be represented as fractions or decimals. In decimal form, it can terminate or repeat. A repeating decimal has a period and a length of period.

Write the fraction as a decimal, then state the period and length of period:

a) $\frac{2}{3} = 0.\overline{6}$ Period: 6 Length of period: 1	b) $\frac{10}{7} = 1.\overline{428571}$ Period: 428571 Length: 6	c) $\frac{5}{12} = 0.41\overline{6}$ Period: 6 Length: 1
---	--	--

$\frac{1}{2} = 0.5$

If the decimal is a terminating decimal, it can be quickly converted to a fraction. (Repeating decimals can be converted, but it can be more complicated and we will not do it here.) The denominator is the place value of the most right digit. The numerator is the number without the decimal. To finish it off, simplify the fraction to lowest terms.

Write the decimal as a fraction in lowest terms:

a) 0.6 $= \frac{6 \div 2}{10 \div 2} = \frac{3}{5}$	b) 1.42 <sup>✓</sup> $= \frac{142 \div 2}{100 \div 2} = \frac{71}{50}$	c) -0.875 $= \frac{-875 \div 5}{1000 \div 5} = \frac{-175 \div 5}{200 \div 5} = \frac{-35 \div 5}{40 \div 5} = -\frac{7}{8}$	d) -3.25 $= \frac{-325 \div 25}{100 \div 25} = -\frac{13}{4}$
--	---	---	--

Put the following numbers in order from lowest to highest:

$$\frac{1}{3}, 0.33, \frac{9}{24}, \sqrt{10}, \frac{3}{10}$$

$$0.\overline{3}, 0.33, 0.375, 3.16, 0.3$$

$$0.3, 0.33, 0.\overline{3}, 0.375, 3.16$$