



13. 14. Given the following triangle, state the six trigonometric ratios for $\angle A$.



$$\begin{aligned} \sin A &= \frac{8}{17} & \csc A &= \frac{17}{8} \\ \cos A &= \frac{15}{17} & \sec A &= \frac{17}{15} \\ \tan A &= \frac{8}{15} & \cot A &= \frac{15}{8} \end{aligned}$$


14. 14. Determine the exact value of $2 \sin^2 60^\circ \times \tan 30^\circ$.



$$\begin{aligned} &2 \sin^2(60) \times \tan(30) \\ &= 2 \left(\frac{\sqrt{3}}{2}\right)^2 \left(\frac{1}{\sqrt{3}}\right) \\ &= 2 \left(\frac{3}{4}\right) \left(\frac{1}{\sqrt{3}}\right) = \frac{6}{4\sqrt{3}} = \frac{3}{2\sqrt{3}} \cdot \frac{\sqrt{3}}{\sqrt{3}} = \frac{3\sqrt{3}}{2(3)} = \frac{\sqrt{3}}{2} \end{aligned}$$


required by some people
acceptable to me
rationalizing the denominator

15. 14. Given $\cos(\theta + 25^\circ) = 0.2237$, solve for θ to the nearest degree. Assume θ is in quadrant 1.




$$\begin{aligned} \cos(\theta + 25^\circ) &= 0.2237 \\ \Rightarrow \theta + 25^\circ &= \cos^{-1}(0.2237) \\ &= 77^\circ \\ \Rightarrow \theta &= 77 - 25 = 52^\circ \end{aligned}$$


16. 17. For each question, draw the angle of rotation and determine the EXACT trig ratio:



$$\sin(150^\circ) = +\frac{1}{2}$$



$$\tan(-315^\circ) = +1$$



$$\sec(210) = -\frac{2}{\sqrt{3}} \left(= -\frac{2\sqrt{3}}{3} \right)$$

$\sec(210^\circ) = \frac{\text{hyp}}{\text{adj}}$
 $\frac{1}{\cos(210^\circ)}$

acceptable.

18. Given the trig ratio determine **both possible** values for θ where $0^\circ \leq \theta \leq 360^\circ$:


17. a) $\sin(\theta) = -\frac{\sqrt{3}}{2}$

b) $\cos(\theta) = \frac{1}{\sqrt{2}}$

c) $\tan(\theta) = -\sqrt{3}$

Q2 or Q4


Q3 or Q4



$\phi = 60^\circ$

$\theta_1 = 180 + \phi = 240^\circ$


$\theta_2 = 360 - \phi = 300^\circ$



$\phi = 45^\circ$

$\theta_1 = 45^\circ$

$\theta_2 = 360 - \phi = 315^\circ$



$\phi = 60^\circ$

$\theta_1 = 180 - \phi = 120^\circ$

$\theta_2 = 360 - \phi = 300^\circ$

19. Given the (inexact!!) trig ratio, determine **both possible** values for θ where $0^\circ \leq \theta \leq 360^\circ$ (round to the nearest degree):


18. a) $\cos(\theta) = -0.3421$

tells us Q2 & Q3

b) $\csc(\theta) = 1.512$

reciprocal of SINE

tells us Q1 & Q2




IGNORE THE NEGATIVE and find ϕ !

$\cos(\phi) = +0.3421$

$\Rightarrow \phi = \cos^{-1}(0.3421)$
 $= 71^\circ$

$\theta_1 = 180 - \phi = 109^\circ$

$\theta_2 = 180 + \phi = 251^\circ$



$\csc(\phi) = 1.512$

$\Rightarrow \sin(\phi) = \frac{1}{1.512} = 0.661$

$\Rightarrow \phi = \sin^{-1}(0.661) = 41^\circ$

$\theta_1 = 41^\circ$

$\theta_2 = 180 - \phi = 139^\circ$

20. Prove the following identity.

$$1 = \frac{(\sin^4 x - \cos^4 x)}{\tan x \sin x \cos x - \cos^2 x}$$

$$\text{RHS} = \frac{\sin^4 x - \cos^4 x}{\tan x \cdot \sin x \cdot \cos x - \cos^2 x}$$

$$= \frac{(\sin^2 x - \cos^2 x)(\sin^2 x + \cos^2 x)}{\tan x \cdot \sin x \cdot \cos x - \cos^2 x}$$

$$= \frac{\sin^2 x \cdot \cos x - \cos^2 x}{\sin^2 x - \cos^2 x}$$

$$= \frac{\sin^2 x - \cos^2 x}{\sin^2 x - \cos^2 x} = 1 = \text{LHS} \quad \square$$

21. Prove the following identity.

$$\tan x \sec x = \frac{\sin^3 x + \cos^2 x \sin x}{\cos^2 x}$$

$$\text{RHS} = \frac{\sin^3 x + \cos^2 x \cdot \sin x}{\cos^2 x}$$

$$= \frac{\sin x (\sin^2 x + \cos^2 x)}{\cos^2 x}$$

$$= \frac{\sin x}{\cos^2 x}$$

(you might see the solution here: $\frac{\sin x}{\cos x} \cdot \frac{1}{\cos x} = \tan x \cdot \sec x$ Boom)

OR switch sides


$$\text{LHS} = \tan(x) \cdot \sec(x)$$

$$= \frac{\sin x}{\cos(x)} \cdot \frac{1}{\cos(x)}$$

$$= \frac{\sin x}{\cos^2 x} = \text{RHS} \quad \square$$

22. A triangular plot of land is enclosed by a fence. One side of the fence is 8.1 m long with an opposite angle of 75° . An adjacent side of the fence is 5.7 m long with an opposite angle of θ .

- a) Make a sketch of the situation.
b) Determine θ to the nearest degree.




$$\frac{\sin \theta}{5.7} = \frac{\sin 75}{8.1}$$

$$\Rightarrow \theta = \sin^{-1} \left(\frac{(5.7)(\sin(75))}{8.1} \right)$$

$$\approx 43^\circ$$

25. The posts of a hockey goal are 2.0 m apart. A player attempts to score by shooting the puck along the ice from a point 7.2 m from one post and 8.6 m from the other. Within what angle θ must the shot be made? Round your answer to the nearest degree.



$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$



$$\Rightarrow \cos A = \frac{8.6^2 + 7.2^2 - 2.0^2}{2(8.6)(7.2)}$$

$$\Rightarrow A = \cos^{-1} \left(\frac{8.6^2 + 7.2^2 - 2.0^2}{2(8.6)(7.2)} \right) \approx 10^\circ$$

\therefore The player needs to shoot within an angle of 10° .


24. Given a triangle with 3 angles that sum to 180° , can the lengths of the sides be determined?

21. No - we need at least one side to determine the \triangle





← Different sized \triangle s with equal angles. No way to calculate the sides.

25. Mary stands on a balcony. Joe is on the left of the balcony looking up at her at an angle of 52° with the ground. Trent is on the right of the balcony looking up at her at an angle of 47° with the ground. If the height, h , is 4 m, how far apart are Joe and Trent standing to the nearest tenth of a metre? Assume the angle the base of the balcony makes between Joe and Trent is 90° .



both of those sides are "common" sides




$$\tan(47^\circ) = \frac{4}{TB} \quad \tan(52^\circ) = \frac{4}{DJ}$$

$$\Rightarrow TB = \frac{4}{\tan(47^\circ)} \quad \Rightarrow DJ = \frac{4}{\tan(52^\circ)}$$

$$= 3.73 \quad = 3.12$$

\therefore Trent and Joe are 4.9 m apart.



By Pythagoras:


$$TJ^2 = 3.73^2 + 3.12^2$$

$$\Rightarrow TJ = \sqrt{\quad}$$

$$= 4.9 \text{ m}$$

26. Doug is looking at a cliff. He determines that the angle of elevation to the top is 54° from where he is at. 50 m away from Doug, Gary estimates the angle between the base of the cliff, himself, and Doug to be 26° while Doug estimates the angle between the base of the cliff, himself, and Gary to be 70° . What is the height, h , of the cliff to the nearest tenth of a metre?

23. need a side in $\triangle CDB$!



we want THIS side (because it is also in $\triangle CDB$)

$\triangle DBG$


Bas! SINE LAW

$$\angle B = 180 - 70 - 26 = 84^\circ$$

$$\frac{DB}{\sin(26)} = \frac{50}{\sin(84)}$$

$$\Rightarrow DB = \frac{50 \cdot \sin(26)}{\sin(84)} = 22.04 \text{ m.}$$

SOH CAH TOA!



$$\tan(54) = \frac{h}{22.04}$$

$$\Rightarrow h = (22.04)(\tan(54)) = 30.3 \text{ m}$$

\therefore The cliff is approximately 30.3 m tall.

I hope this has helped!