





Chapter

4

Polynomial Equations and Inequalities

► GOALS

You will be able to


- Determine the roots of polynomial equations, with and without technology
- Solve polynomial inequalities, with and without technology
- Solve problems involving polynomial function models

? If a polynomial function models the height of the wake boarder above the surface of the water, how could you use the function to determine when he is above a given height or how quickly he is descending at any given time?

Study Aid

- For help, see the Review of Essential Skills found at the Nelson Advanced Functions website.

Question	Appendix/Lesson
1, 4, 5	R-6
2	R-3, 3.6
3	3.3
6, 7	2.5



SKILLS AND CONCEPTS You Need

1. Solve the following linear equations.

a) $5x - 7 = -3x + 17$ c) $2(3x - 5) = -4(3x - 2)$
 b) $12x - 9 - 6x = 5 + 3x + 1$ d) $\frac{2x + 5}{3} = 7 - \frac{x}{4}$

2. Factor the following expressions.

a) $x^3 + x^2 - 30x$ c) $24x^4 + 81x$
 b) $x^3 - 64$ d) $2x^3 + 7x^2 - 18x - 63$

3. Sketch a graph of each of the following functions.


a) $y = (x - 2)(x + 3)(x - 4)$ b) $y = 2(x + 6)^3 - 10$

4. Given the graph of the function shown, determine the roots of the equation $x^2 - 7x + 10 = 0$.

5. Determine the roots of each of the following quadratic equations.

a) $2x^2 = 18$ c) $6x^2 = 11x + 10$
 b) $x^2 + 8x - 20 = 0$ d) $x(x + 3) = 3 - 5x - x^2$

6. The graph below shows Erika's walk in front of a motion sensor.



- a) In which time interval is she walking the fastest? Explain.
 b) Calculate the speeds at which she walks on the intervals $t \in (0, 3)$ and $t \in (3, 7)$.
 c) Is she moving away or toward the motion sensor? How do you know?
7. A T-ball player hits a baseball from a tee that is 0.5 m tall. The height of the ball is modelled by $h(t) = -5t^2 + 9.75t + 0.5$, where h is the height in metres at t seconds.
- a) How long is the ball in the air?
 b) Determine the average rate of change in the ball's height during the first second of flight.
 c) Estimate the instantaneous rate of change in the ball's height when it hits the ground.

8. Copy and complete the anticipation guide in your notes.

Statement	Agree	Disagree	Justification
The quadratic formula can only be used when solving a quadratic equation.			
Cubic equations always have three real roots.			
The graph of a cubic function always passes through all four quadrants.			
The graphs of all polynomial functions must pass through at least two quadrants.			
The expression $x^2 > 4$ is only true if $x > 2$.			
If you know the instantaneous rates of change for a function at $x = 2$ and $x = 3$, you can predict fairly well what the function looks like in between.			

APPLYING What You Know

Modelling a Situation with a Polynomial Function

Shown is a picture of the Gateway Arch located in St. Louis, Missouri, U.S.A. The arch is about 192 m wide and 192 m tall.

The city of St. Louis would like to hang a banner from the arch for their New Year celebrations. They have determined that the banner should be suspended from a horizontal cable that spans the arch 175 m off the ground to ensure optimal viewing around the city.



- ? Assuming that the arch is parabolic in shape, how long should the cable be?**
- Draw a sketch showing the arch on a coordinate grid using an appropriate scale.
 - Determine a quadratic function that could model the inside of the arch using vertex or factored form.
 - How did you predict what the sign of the leading coefficient in the function would be? Explain.
 - Use your model to determine the length of the cable needed to support the banner.

4.1

Solving Polynomial Equations

YOU WILL NEED

- graphing calculator or graphing software



GOAL

Solve polynomial equations using a variety of strategies.

LEARN ABOUT the Math

Amelia's family is planning to build another silo for grain storage, identical to those they have on their farm. The cylindrical portion of those they currently have is 15 m tall, and the silo's total volume is $684\pi \text{ m}^3$.


? What are the possible values for the radius of the new silo?

EXAMPLE 1

Representing a problem with a polynomial model

Determine possible values for the radius of the silo.

Solution



Draw a diagram to represent the silo.
In this case, the height must be 15 m.

$$V = V_{\text{cylinder}} + V_{\text{hemisphere}}$$

$$V = \pi r^2 h + \frac{1}{2} \left(\frac{4}{3} \pi r^3 \right)$$

Determine a **polynomial equation** for the volume of the silo using the formula for the volume of a cylinder and a hemisphere.

$$684\pi = \pi r^2 (15) + \frac{2}{3} \pi r^3$$

$$684\pi = 15\pi r^2 + \frac{2}{3} \pi r^3$$

Substitute the given values for the volume and the height into the formula and simplify the equation.

$$0 = 15\pi r^2 + \frac{2}{3} \pi r^3 - 684\pi$$

$$0 = \frac{\pi}{3} (45r^2 + 2r^3 - 2052)$$

Divide out the common factor of $\frac{\pi}{3}$, then divide both sides of the equation by this value.

polynomial equation

an equation in which one polynomial expression is set equal to another (e.g., $x^3 - 5x^2 = 4x - 3$, or $5x^4 - 3x^3 + x^2 - 6x = 9$)

$$0 = 45r^2 + 2r^3 - 2052$$

$$0 = 2r^3 + 45r^2 - 2052 \quad \leftarrow \begin{array}{l} \text{Rewrite the polynomial in} \\ \text{descending order of degree.} \end{array}$$

Let $f(r) = 2r^3 + 45r^2 - 2052 \quad \leftarrow \begin{array}{l} \text{Solve the equation by} \\ \text{factoring the polynomial.} \end{array}$

$$f(2) = 2(2)^3 + 45(2)^2 - 2052$$

$$= -1856$$

$$f(3) = 2(3)^3 + 45(3)^2 - 2052$$

$$= -1593 \quad \leftarrow \begin{array}{l} \text{Since the roots of the equation} \\ \text{are the x-intercepts of the} \\ \text{related function, use the factor} \\ \text{theorem to determine one} \\ \text{factor.} \end{array}$$

$$f(6) = 2(6)^3 + 45(6)^2 - 2052$$

$$= 0$$

By the factor theorem, $(r - 6)$ is a factor of $f(r)$.

$$(2r^3 + 45r^2 - 2052) \div (r - 6)$$

$$\begin{array}{r|rrrr} 6 & 2 & 45 & 0 & -2052 \\ & \downarrow & 12 & 342 & 2052 \\ \hline & 2 & 57 & 342 & 0 \end{array} \quad \leftarrow \begin{array}{l} \text{Use synthetic division to divide} \\ f(r) \text{ by } (r - 6) \text{ to determine} \\ \text{the other factor.} \end{array}$$

$$(r - 6)(2r^2 + 57r + 342) = 0$$

$$r - 6 = 0$$

$$r = 6 \text{ is one solution} \quad \leftarrow \begin{array}{l} \text{Set each factor equal to zero} \\ \text{and solve.} \end{array}$$

$$2r^2 + 57r + 342 = 0$$

$$a = 2, b = 57, c = 342$$

$$r = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$r = \frac{-57 \pm \sqrt{(57)^2 - 4(2)(342)}}{2(2)}$$

$$r = \frac{-57 \pm \sqrt{513}}{4}$$

$$r = \frac{-57 + \sqrt{513}}{4} \text{ or } r = \frac{-57 - \sqrt{513}}{4}$$

$$r \doteq -8.6 \quad r \doteq -19.9$$

The silo must have a radius of 6 m. $\leftarrow \begin{array}{l} \text{The radius cannot be negative,} \\ \text{and so only the positive root} \\ \text{can be the radius of the silo.} \end{array}$

Reflecting

- How could you verify the solutions you found, with and without using a graphing calculator?
- What restriction was placed on the variable in the polynomial equation? Explain why this was necessary.
- Do you think it is possible to solve all cubic and quartic equations using an algebraic strategy involving factoring? Explain.

APPLY the Math

EXAMPLE 2 Selecting a strategy to solve a cubic equation

Solve $4x^3 - 12x^2 - x + 3 = 0$.

Solution A: Using the factor theorem

Let $f(x) = 4x^3 - 12x^2 - x + 3$

Possible values for x where $f(x) = 0$:

$$\pm 1, \pm 3, \pm \frac{1}{2}, \pm \frac{3}{2}, \pm \frac{1}{4}, \pm \frac{3}{4}$$

$$\begin{aligned} f(1) &= 4(1)^3 - 12(1)^2 - (1) + 3 \\ &= -6 \end{aligned}$$

$$\begin{aligned} f(-1) &= 4(-1)^3 - 12(-1)^2 - (-1) + 3 \\ &= -12 \end{aligned}$$

$$\begin{aligned} f(3) &= 4(3)^3 - 12(3)^2 - (3) + 3 \\ &= 0 \end{aligned}$$

By the factor theorem, $(x - 3)$ is a factor of $f(x)$.

$$(4x^3 - 12x^2 - x + 3) \div (x - 3)$$

$$\begin{array}{r} 4x^2 + 0x - 1 \\ x - 3 \overline{) 4x^3 - 12x^2 - x + 3} \\ \underline{4x^3 - 12x^2} \\ 0x^2 - x \\ \underline{0x^2 - 0x} \\ -x + 3 \\ \underline{-x + 3} \\ 0 \end{array}$$

$$(x - 3)(4x^2 - 1) = 0$$

Use the factor theorem and the related polynomial function to determine one factor of the equation. Numbers that could make $f(x) = 0$ are of the form $\frac{p}{q}$, where p is a factor of the constant term 3 and q is a factor of the leading coefficient 4.

Divide $f(x)$ by $(x - 3)$ to find the second factor using either long or synthetic division.

The quotient $4x^2 - 1$ is a difference of squares. Factor this.



$$(x - 3)(2x - 1)(2x + 1) = 0$$

$$x - 3 = 0 \text{ or } 2x - 1 = 0 \text{ or } 2x + 1 = 0 \quad \leftarrow$$

$$x = 3 \qquad 2x = 1 \qquad 2x = -1$$

$$x = \frac{1}{2} \qquad x = -\frac{1}{2}$$

Set each of the factors equal to zero to solve.

Check:

$$x = \frac{1}{2}$$

$\begin{aligned} \text{LS} \\ 4x^3 - 12x^2 - x + 3 \\ = 4\left(\frac{1}{2}\right)^3 - 12\left(\frac{1}{2}\right)^2 \\ - \left(\frac{1}{2}\right) + 3 \\ = \frac{1}{2} - 3 - \frac{1}{2} + 3 \\ = 0 \end{aligned}$	RS	$\begin{aligned} 0 \\ \text{LS} = \text{RS} \checkmark \end{aligned}$
---	----	---

$$x = -\frac{1}{2}$$

$\begin{aligned} \text{LS} \\ 4x^3 - 12x^2 - x + 3 \\ = 4\left(-\frac{1}{2}\right)^3 - 12\left(-\frac{1}{2}\right)^2 \\ - \left(-\frac{1}{2}\right) + 3 \\ = -\frac{1}{2} - 3 + \frac{1}{2} + 3 \\ = 0 \end{aligned}$	RS	$\begin{aligned} 0 \\ \text{LS} = \text{RS} \checkmark \end{aligned}$
---	----	---

Verify the solutions by substitution.

You only need to check $x = \frac{1}{2}$ and $-\frac{1}{2}$ since $x = 3$ was obtained using substitution.

The solutions to $4x^3 - 12x^2 - x + 3 = 0$ are $x = -\frac{1}{2}$, $x = \frac{1}{2}$, and $x = 3$.

Solution B: Factoring by grouping

$$4x^3 - 12x^2 - x + 3 = 0$$

$$4x^2(x - 3) - 1(x - 3) = 0 \quad \leftarrow$$

$$(x - 3)(4x^2 - 1) = 0$$

$$(x - 3)(2x - 1)(2x + 1) = 0$$

$$x - 3 = 0 \text{ or } 2x - 1 = 0 \text{ or } 2x + 1 = 0 \quad \leftarrow$$


$$x = 3 \qquad 2x = 1 \text{ or } \qquad 2x = -1$$

$$x = \frac{1}{2} \qquad x = -\frac{1}{2}$$

The first two terms and the last two terms have a common factor, so you can factor by grouping.

Set each of the factors equal to zero to solve.





Verify the solutions, by graphing the corresponding polynomial function and determining its zeros.

The solutions to $4x^3 - 12x^2 - x + 3 = 0$ are $x = -\frac{1}{2}$, $x = \frac{1}{2}$, and $x = 3$.

EXAMPLE 3 Selecting tools to solve a question involving modelling



The paths of two orcas playing in the ocean were recorded by some oceanographers. The first orca's path could be modelled by the equation $h(t) = 2t^4 - 17t^3 + 27t^2 - 252t + 232$, and the second by $h(t) = 20t^3 - 200t^2 + 300t - 200$, where h is their height above/below the water's surface in centimetres and t is the time during the first 8 s of play. Over this 8-second period, at what times were the two orcas at the same height or depth?

Solution

$$2t^4 - 17t^3 + 27t^2 - 252t + 232 = 20t^3 - 200t^2 + 300t - 200$$

$$2t^4 - 37t^3 + 227t^2 - 552t + 432 = 0$$

Since you are solving for the time when the heights or depths are the same, set the two equations equal to each other and use inverse operations to make the right side of the equation equal to zero.

$$\text{Let } f(t) = 2t^4 - 37t^3 + 227t^2 - 552t + 432$$

Solve the equation by factoring.



Some possible values for t where $f(t) = 0$:

$$\pm 1, \pm 2, \pm 3, \pm 4, \pm 6, \pm 8, \pm 9$$

$$f(1) = 72$$

$$f(2) = -28$$

$$f(3) = -18$$

$$f(4) = 0$$

By the factor theorem, $(t - 4)$ is a factor of $f(t)$.

$$(2t^4 - 37t^3 + 227t^2 - 552t + 432) \div (t - 4)$$

$$\begin{array}{r|rrrrr} 4 & 2 & -37 & 227 & -552 & 432 \\ & \downarrow & & & & \\ & & 8 & -116 & 444 & -432 \\ \hline & 2 & -29 & 111 & -108 & 0 \end{array}$$

$$f(t) = (t - 4)(2t^3 - 29t^2 + 111t - 108)$$

$$f(6) = -108$$

$$f(8) = -208$$

$$f(9) = 0$$

By the factor theorem, $(t - 9)$ is a factor of $f(t)$.

$$(2t^3 - 29t^2 + 111t - 108) \div (t - 9)$$

$$\begin{array}{r|rrrr} 9 & 2 & -29 & 111 & -108 \\ & \downarrow & & & \\ & & 18 & -99 & 108 \\ \hline & 2 & -11 & 12 & 0 \end{array}$$

$$f(t) = (t - 4)(t - 9)(2t^2 - 11t + 12)$$

$$= (t - 4)(t - 9)(2t - 3)(t - 4)$$

$$= (t - 4)^2(t - 9)(2t - 3)$$

$$(t - 4)^2 = 0, t - 9 = 0, \text{ or } 2t - 3 = 0$$

$$t = 4 \qquad t = 9 \qquad t = 1.5$$

The solutions that are valid on the given domain are $t = 1.5$ and $t = 4$.

Use the factor theorem to determine one factor of $f(t)$. Since the question specifies that the time is within the first 10 s, you only need to consider values of $\frac{p}{q}$ between 0 and 10. In this case, consider just the factors of 432.

Divide $f(t)$ by $(t - 4)$ to determine the second factor.

Since the second factor is cubic, you must continue looking for more zeros using the factor theorem. It is not necessary to recheck the values that were used in an earlier step, so carry on with the other possibilities.



Divide the cubic polynomial by $(t - 9)$ to determine the other factor.

Factor the quadratic.

Set each factor equal to zero and solve.

The polynomial functions given only model the orca's movement between 0 s and 8 s, so the solution $t = 9$ is inadmissible.





Graph both functions and adjust the window settings to determine the points of intersection to verify the solutions.

The orcas were at the same depth after 1.5 s and 4 s on the interval between 0 s and 8 s.

EXAMPLE 4

Selecting a strategy to solve a polynomial equation that is unfactorable

Solve each of the following.

a) $x^4 + 5x^2 = -1$ b) $x^3 - 2x + 3 = 0$

Solution

a) $x^4 + 5x^2 = -1$
 $x^4 + 5x^2 + 1 = 0$


Add 1 to both sides of the equation to make the right side of the equation equal to zero.

Let $f(x) = x^4 + 5x^2 + 1$
 $f(1) = (1)^4 + 5(1)^2 + 1 = 7$
 $f(-1) = (-1)^4 + 5(-1)^2 + 1 = 7$

If the equation is factorable, then either $f(1)$ or $f(-1)$ should give a value of 0.
 The polynomial in this equation cannot be factored.

The function $f(x) = x^4 + 5x^2 + 1$ has an even degree and a positive leading coefficient, so its end behaviours are the same. In this case, as $x \rightarrow \pm\infty$, $y \rightarrow \infty$. This function has a degree of 4, so it could have 4, 3, 2, 1, or 0 x -intercepts.

Use the corresponding polynomial function to visualize the graph and determine the possible number of zeros.



Graph the function on a graphing calculator to determine its zeros.

Based on the graph and the function's end behaviours, it never crosses the x -axis, so it has no zeros. As a result, the equation $x^4 + 5x^2 = -1$ has no solutions.

b) $x^3 - 2x + 3 = 0$

Let $f(x) = x^3 - 2x + 3$

$$f(1) = (1)^3 - 2(1) + 3 = 2$$

$$f(-1) = (-1)^3 - 2(-1) + 3 = 4$$

$$f(3) = (3)^3 - 2(3) + 3 = 24$$


$$f(-3) = (-3)^3 - 2(-3) + 3 = -18$$

If the equation is factorable, then either $f(1)$, $f(-1)$, $f(3)$, or $f(-3)$ should equal 0.

The polynomial in this equation cannot be factored.

The function $f(x) = x^3 - 2x^2 + 3$ has an odd degree and a positive leading coefficient, so its end behaviours are opposite. In this case, as $x \rightarrow \infty$, $y \rightarrow \infty$, and as $x \rightarrow -\infty$, $y \rightarrow -\infty$. This function has a degree of 3, so it could have 3, 2, or 1 x -intercepts.

Use the corresponding polynomial function to visualize the graph and determine the possible number of zeros.



Graph the function on a graphing calculator to determine its zeros.

Based on the graph and the function's end behaviours, it crosses the x -axis only once. The solution is $x \doteq -1.89$.

In Summary

Key Idea

- The solutions to a polynomial equation $f(x) = 0$ are the zeros of the corresponding polynomial function, $y = f(x)$.

Need to Know

- Polynomial equations can be solved using a variety of strategies:
 - algebraically using a factoring strategy
 - graphically using a table of values, transformations, or a graphing calculator
- Only some polynomial equations can be solved by factoring, since not all polynomials are factorable. In these cases, graphing technology must be used.
- When solving problems using polynomial models, it may be necessary to ignore the solutions that are outside the domain defined by the conditions of the problem.

CHECK Your Understanding

- State the zeros of the following functions.
 - $y = 2x(x - 1)(x + 2)(x - 2)$
 - $y = 5(2x + 3)(4x - 5)(x + 7)$
 - $y = 2(x - 3)^2(x + 5)(x - 4)$
 - $y = (x + 6)^3(2x - 5)$
 - $y = -5x(x^2 - 9)$
 - $y = (x + 5)(x^2 - 4x - 12)$
- Solve each of the following equations by factoring. Verify your solutions using graphing technology.
 - $3x^3 = 27x$
 - $4x^4 = 24x^2 + 108$
 - $3x^4 + 5x^3 - 12x^2 - 20x = 0$
 - $10x^3 + 26x^2 - 12x = 0$
 - $2x^3 + 162 = 0$
 - $2x^4 = 48x^2$
- Determine the zeros of the function $y = 2x^3 - 17x^2 + 23x + 42$.
 - Write the polynomial equation whose roots are the zeros of the function in part a).
- Explain how you can solve $x^3 + 12x^2 + 21x - 4 = x^4 - 2x^3 - 13x^2 - 4$ using two different strategies.
- Determine the zeros of the function $f(x) = 2x^4 - 11x^3 - 37x^2 + 156x$ algebraically. Verify your solution using graphing technology.

PRACTISING

- State the zeros of the following functions.
 - $f(x) = x(x - 2)^2(x + 5)$
 - $f(x) = (x^3 + 1)(x - 17)$
 - $f(x) = (x^2 + 36)(8x - 16)$
 - $f(x) = -3x^3(2x + 4)(x^2 - 25)$
 - $f(x) = (x^2 - x - 12)(3x)$
 - $f(x) = (x + 1)(x^2 + 2x + 1)$
- Determine the roots algebraically by factoring.
 - $x^3 - 8x^2 - 3x + 90 = 0$
 - $x^4 + 9x^3 + 21x^2 - x - 30 = 0$
 - $2x^3 - 5x^2 - 4x + 3 = 0$
 - $2x^3 + 3x^2 = 5x + 6$
 - $4x^4 - 4x^3 - 51x^2 + 106x = 40$
 - $12x^3 - 44x^2 = -49x + 15$

8. Use graphing technology to find the real roots to two decimal places.

a) $x^3 - 7x + 6 = 0$

d) $x^5 + x^4 = 5x^3 - x^2 + 6x$

b) $x^4 - 5x^3 - 17x^2 + 3x + 18 = 0$

e) $105x^3 = 344x^2 - 69x - 378$

c) $3x^3 - 2x^2 + 16 = x^4 + 16x$

f) $21x^3 - 58x^2 + 10 = -18x^4 - 51x$

9. Solve each of the following equations.


K a) $x^3 - 6x^2 - x + 30 = 0$

b) $9x^4 - 42x^3 + 64x^2 - 32x = 0$

c) $6x^4 - 13x^3 - 29x^2 + 52x = -20$

d) $x^4 - 6x^3 + 10x^2 - 2x = x^2 - 2x$

10. An open-topped box can be created by cutting congruent squares from each of the four corners of a piece of cardboard that has dimensions of 20 cm by 30 cm and folding up the sides. Determine the dimensions of the squares that must be cut to create a box with a volume of 1008 cm^3 .



11. The Sickle-Lichti family members are very competitive card players.

A They keep score using a complicated system that incorporates positives and negatives. Maya's score for the last game night could be modelled by the function $S(x) = x(x - 4)(x - 6)$, $x < 10$, $x \in \mathbf{W}$, where x represents the game number.

- After which game was Maya's score equal to zero?
- After which game was Maya's score -5 ?
- After which game was Maya's score 16?
- Draw a sketch of the graph of $S(x)$ if $x \in \mathbf{R}$. Explain why this graph is *not* a good model to represent Maya's score during this game night.

12. The function $s(t) = -\frac{1}{2}gt^2 + v_0t + s_0$ can be used to calculate s , the height above a planet's surface in metres, where g is the acceleration due to gravity, t is the time in seconds, v_0 is the initial velocity in metres per second, and s_0 is the initial height in metres. The acceleration due to gravity on Mars is $g = -3.92 \text{ m/s}^2$. Find, to two decimal places, how long it takes an object to hit the surface of Mars if the object is dropped from 1000 m above the surface.

13. The distance of a ship from its harbour is modelled by the function $d(t) = -3t^3 + 3t^2 + 18t$, where t is the time elapsed in hours since departure from the harbour.
- Factor the time function.
 - When does the ship return to the harbour?
 - There is another zero of $d(t)$. What is it, and why is it not relevant to the problem?
 - Draw a sketch of the function where $0 \leq t \leq 3$.
 - Estimate the time that the ship begins its return trip back to the harbour.
14. During a normal 5 s respiratory cycle in which a person inhales and then exhales, the volume of air in a person's lungs can be modelled by $V(t) = 0.027t^3 - 0.27t^2 + 0.675t$, where the volume, V , is measured in litres at t seconds.
- What restriction(s) must be placed on t ?
 - If asked, "How many seconds have passed if the volume of air in a person's lungs is 0.25 L?" would you answer this question algebraically or by using graphing technology? Justify your decision.
 - Solve the problem in part b).
15. Explain why the following polynomial equation has no real solutions:
 $0 = 5x^8 + 10x^6 + 7x^4 + 18x^2 + 132$
16. Determine algebraically where the cubic polynomial function that has **T** zeros at 2, 3, and -5 and passes through the point $(4, 36)$ has a value of 120.
17. For each strategy below, create a cubic or quartic equation you might **C** solve by using that strategy (the same equation could be used more than once). Explain why you picked the equation you did.
- | | |
|-----------------------|-------------------------------|
| a) factor theorem | d) quadratic formula |
| b) common factor | e) difference or sum of cubes |
| c) factor by grouping | f) graphing technology |

Extending

18. a) It is possible that a polynomial equation of degree 4 can have no real roots. Create such a polynomial equation and explain why it cannot have any real roots.
- b) Explain why a degree 5 polynomial equation must have at least one real root.
19. The factor theorem only deals with rational zeros. Create a polynomial of degree 5 that has no rational zeros. Explain why your polynomial has no rational zero but has at least one irrational zero.

4.2

Solving Linear Inequalities

GOAL

Solve linear inequalities.

YOU WILL NEED

- graphing calculator or graphing software

LEARN ABOUT the Math

In mathematics, you must be able to represent intervals and identify smaller sections of a relation or a set of numbers. You have used the following inequality symbols:

$>$ greater than $<$ less than
 \geq greater than or equal to \leq less than or equal to


When you write one of these symbols between two or more linear expressions, the result is called a **linear inequality**.

To solve an inequality, you have to find all the possible values of the variable that satisfy the inequality.

For example, $x = 2$ satisfies $3x - 1 < 8$, but so do $x = 2.9$, $x = -1$, and $x = -5$.



In fact, every real number less than 3 results in a number smaller than 8. So all real numbers less than 3 satisfy this inequality. The thicker part of the number line below represents this solution.



The solution to $3x - 1 < 8$ can be written in set notation as $\{x \in \mathbf{R} \mid x < 3\}$ or in interval notation $x \in (-\infty, 3)$.

? How can you determine algebraically the solution set to a linear inequality like $3x - 1 < 8$?

linear inequality

an inequality that contains an algebraic expression of degree 1 (e.g., $5x + 3 > 6x - 2$)

Communication Tip

To show that a number is not included in the solution set, use an open dot at this value. A solid dot shows that this value is included in the solution set.

EXAMPLE 1 Selecting an inverse operation strategy to solve a linear inequality

Solve the linear inequality $3x - 1 < 8$.

Solution

$$3x - 1 < 8$$

$$3x - 1 + 1 < 8 + 1$$


$$3x < 9$$

$$\frac{3x}{3} < \frac{9}{3}$$

$$x < 3$$

Treat the inequality like a linear equation and use inverse operations to isolate x . Add 1 to both sides of the inequality and simplify.

Divide both sides of the inequality by 3.





A number line helps visualize the solution.

Check $x = 0$.


LS	RS
$3x - 1$	8
$= 3(0) - 1$	
$= -1$	

LS < RS

Choose a value for x that is less than 3 to verify that any number in the solution set satisfies the original inequality.



You can also verify the solution set using a graphing calculator. Graph each side of the inequality as a function.



The solution set is $\{x \in \mathbf{R} \mid x < 3\}$ or in interval notation $x \in (-\infty, 3)$.

The y -values on the line $y = 3x - 1$ that are less than 8 are found on all points that lie on the line below the horizontal line $y = 8$. This happens when x is smaller than 3.

Reflecting

- A. How was solving a linear inequality like solving a linear equation?
How was it different?
- B. When checking the solution to an inequality, why is it not necessary for the left side to equal the right side?
- C. Why do most linear equations have only one solution, but linear inequalities have many?

APPLY the Math

EXAMPLE 2

Using reasoning to determine which operations preserve the truth of a linear inequality

Can you add, subtract, multiply, or divide both sides of an inequality by a non-zero value and still have a valid inequality?

Solution

$$4 < 8$$

$$4 + 5 < 8 + 5$$

$$9 < 13$$

The result is still true.

$$4 + (-5) < 8 + (-5)$$

$$-1 < 3$$

The result is still true.

$$4 - 10 < 8 - 10$$

$$-6 < -2$$

The result is still true.

$$4 - (-3) < 8 - (-3)$$

$$7 < 11$$

The result is still true.

$$4(6) < 8(6)$$

$$24 < 48$$

The result is still true.

Write a true inequality using two numbers. Add the same positive quantity to both sides.

Add the same negative quantity to both sides of the initial inequality.

Subtract the same positive quantity from both sides of the initial inequality.

Subtract the same negative quantity from both sides of the initial inequality.

Multiply by the same positive quantity on both sides of the initial inequality.



$$4(-2) < 8(-2)$$

$$-8 < -16$$

Multiply by the same negative quantity on both sides of the initial inequality.

The result is false.
In this case, $-8 > -16$.

$$4 \div 2 < 8 \div 2$$

$$2 < 4$$

Divide by the same positive quantity on both sides of the initial inequality.

The result is still true.

$$4 \div (-2) < 8 \div (-2)$$

$$-2 < -4$$

Divide by the same negative quantity on both sides of the initial inequality.

The result is false.
In this case, $-2 > -4$.

Most of the operations preserve the validity of the inequality. The exception occurs when both sides are multiplied or divided by a negative number. In these two cases, reversing the inequality sign preserves the validity.

Since algebraic expressions represent numbers, this conclusion applies to linear inequalities that contain variables.

EXAMPLE 3

Reflecting to verify a solution

Solve the inequality $35 - 2x \geq 20$.

Solution

$$35 - 2x \geq 20$$

$$-2x \geq 20 - 35$$

$$-2x \geq -15$$

Use inverse operations to isolate x . Subtract 35 from both sides and simplify.

$$\frac{-2x}{-2} \leq \frac{-15}{-2}$$

$$x \leq 7.5$$

Divide both sides by -2 . Since the division involves a negative number, reverse the inequality sign.



Represent the solution on a number line.
A solid dot is placed on 7.5 since this number is included in the solution set.

If $x = 5$, then

LS	RS
$35 - 2x$	20
$= 35 - 2(5)$	
$= 35 - 10$	
$= 25$	

$LS \geq RS$: This is the desired outcome.

Test a value less than 7.5 and a value greater than 7.5 to verify the solution. Since $x = 5$ makes the inequality true and $x = 8$ makes the inequality false, the solution is correct.

If $x = 8$, then

LS	RS
$35 - 2x$	20
$= 35 - 2(8)$	
$= 35 - 16$	
$= 19$	

$LS \leq RS$: This is not the desired outcome.

The solution set is $\{x \in \mathbf{R} \mid x \leq 7.5\}$ or in interval notation $x \in (-\infty, 7.5]$.

The value of 7.5 makes both sides equal. Since the inequality sign includes an equal sign, 7.5 must be part of the solution set.

EXAMPLE 4

Connecting the process of solving a double inequality to solving a linear inequality

Solve the inequality $30 \leq 3(2x + 4) - 2(x + 1) \leq 46$.

Solution

$$30 \leq 3(2x + 4) - 2(x + 1) \leq 46$$

This is a combination of two inequalities:
 $30 \leq 3(2x + 4) - 2(x + 1)$ and
 $3(2x + 4) - 2(x + 1) \leq 46$.
 A valid solution must satisfy both inequalities.
 Expand using the distributive property and simplify.

$$30 \leq 6x + 12 - 2x - 2 \leq 46$$

$$30 \leq 4x + 10 \leq 46$$

$$30 - 10 \leq 4x + 10 - 10 \leq 46 - 10$$

$$20 \leq 4x \leq 36$$

Subtract 10 from all three parts of the inequality.

$$\frac{20}{4} \leq \frac{4x}{4} \leq \frac{36}{4}$$

$$5 \leq x \leq 9$$





The solution using interval notation is


$$x \in [5, 9]$$

Divide all parts of the inequality by 4.

A number line helps to visualize the solution. Solid dots are placed on 5 and 9 since these numbers are included in the solution set.



To verify the solution, graph the functions that correspond to all three parts of the inequality.



The x -values that satisfy the inequality are the x -coordinates of points on the diagonal line defined by $y = 3(2x + 4) - 2(x + 1)$ whose y -values are bounded by 30 and 46.

In Summary

Key Idea

- You can solve a linear inequality using inverse operations in much the same way you solve linear equations.

Need to Know

- If you multiply or divide an inequality by a negative number, you must reverse the inequality sign.
- Most linear equations have only one solution, whereas linear inequalities have many solutions.
- A number line can help you visualize the solution set to an inequality. A solid dot is used to indicate that a number is included in the solution set, whereas an open dot indicates that a number is excluded.

CHECK Your Understanding

- Solve the following inequalities graphically. Express your answer using set notation.

a) $3x - 1 \leq 11$	d) $3(2x + 4) \geq 2x$
b) $-x + 5 > -2$	e) $-2(1 - 2x) < 5x + 8$
c) $x - 2 > 3x + 8$	f) $\frac{6x + 8}{5} \leq 2x - 4$
- Solve the following inequalities algebraically. Express your answer using interval notation.

a) $2x - 5 \leq 4x + 1$	d) $2x + 1 \leq 5x - 2$
b) $2(x + 3) < -(x - 4)$	e) $-x + 1 > x + 1$
c) $\frac{2x + 3}{3} \leq x - 5$	f) $\frac{x + 4}{2} \geq \frac{x - 2}{4}$
- Solve the double inequality $3 \leq 2x + 5 < 17$ algebraically and illustrate your solution on a number line.
- For each of the following inequalities, determine whether $x = 2$ is contained in the solution set.

a) $x > -1$	d) $5x + 3 \leq -3x + 1$
b) $5x - 4 > 3x + 2$	e) $x - 2 \leq 3x + 4 \leq x + 14$
c) $4(3x - 5) \geq 6x$	f) $33 < -10x + 3 < 54$

PRACTISING

- Solve the following algebraically. Verify your results graphically.

K a) $2x - 1 \leq 13$	d) $5(x - 3) \geq 2x$
b) $-2x - 1 > -1$	e) $-4(5 - 3x) < 2(3x + 8)$
c) $2x - 8 > 4x + 12$	f) $\frac{x - 2}{3} \leq 2x - 3$
- For the following inequalities, determine if 0 is a number in the solution set.

a) $3x \leq 4x + 1$	d) $3x \leq x + 1 \leq x - 1$
b) $-6x < x + 4 < 12$	e) $x(2x - 1) \leq x + 7$
c) $-x + 1 > x + 12$	f) $x + 6 < (x + 2)(5x + 3)$
- Solve the following inequalities algebraically.


a) $-5 < 2x + 7 < 11$	d) $0 \leq -2(x + 4) \leq 6$
b) $11 < 3x - 1 < 23$	e) $59 < 7x + 10 < 73$
c) $-1 \leq -x + 9 \leq 13$	f) $18 \leq -12(x - 1) \leq 48$

8. a) Create a linear inequality, with both constant and linear terms on each side, for which the solution is $x > 4$.
 b) Create a linear inequality, with both constant and linear terms on each side, for which the solution is $x \leq \frac{3}{2}$.

9. The following number line shows the solution to a double inequality.



- a) Write the solution using set notation.
 b) Create a double inequality for which this is the solution set.
10. Which of the following inequalities has a solution. Explain.
 $x - 3 < 3 - x < x - 5$ or $x - 3 > 3 - x > x - 5$
11. Consider the following graph.



- a) Write an inequality that is modelled by the graph.
 b) Find the solution by examining the graph.
 c) Confirm the solution by solving your inequality algebraically.
12. The relationship between Celsius and Fahrenheit is represented by $C = \frac{5}{9}(F - 32)$. In order to be comfortable, but also economical, the temperature in your house should be between 18°C and 22°C .
 a) Write this statement as a double linear inequality.
 b) Solve the inequality to determine the temperature range in degrees Fahrenheit.
13. Some volunteers are making long distance phone calls to raise money for a charity. The calls are billed at the rate of \$0.50 for the first 3 min and \$0.10/min for each additional minute or part thereof. If each call cannot cost more than \$2.00, how long can each volunteer talk to a prospective donor?

14. a) Find the equation that allows for the conversion of Celsius to Fahrenheit by solving the relation given in question 12 for F.
 b) For what values of C is the Fahrenheit temperature greater than the equivalent Celsius temperature?
15. The inequality $|2x - 1| < 7$ can be expressed as a double inequality.
T a) Depict the inequality graphically.
 b) Use your graph to solve the inequality.
16. Will the solution to a double inequality always have an upper and lower limit? Explain.
C

Extending

17. Some inequalities are very difficult to solve algebraically. Other methods, however, can be very helpful in solving such problems. Consider the inequality $2^x - 3 < x + 1$.
 a) Explain why solving the inequality might be very difficult to do algebraically.
 b) Describe an alternative method that could work, and use it to solve the inequality.
18. Some operations result in switching the direction of the inequality when done to both sides, but others result in maintaining the direction. For instance, if you add a constant to both sides, the direction is maintained, whereas multiplying both sides by a negative constant causes the sign to switch. For each of the following, determine if the inequality direction should be maintained, should switch, or if it sometimes switches and sometimes is maintained.
 a) cubing both sides
 b) squaring both sides
 c) making each side the exponent with 2 as the base, i.e., $3 < 5$, so $2^3 < 2^5$
 d) making each side the exponent with 0.5 as the base
 e) taking the reciprocal of both sides
 f) rounding both sides up to the nearest integer
 g) taking the square root of both sides
19. Solve each of the following, $x \in \mathbf{R}$. Express your answers using both set and interval notation and graph the solution set on a number line.
 a) $x^2 < 4$ c) $|2x + 2| < 8$
 b) $4x^2 + 5 \geq 41$ d) $-3x^3 \geq 81$

FREQUENTLY ASKED Questions

Study Aid

- See Lesson 4.1, Examples 1 to 4.
- Try Mid-Chapter Review Questions 1, 2, and 3.

Q: How can you solve a polynomial equation?

A1: You can use an algebraic strategy using the corresponding polynomial function, the factor theorem, and division to factor the polynomial. Set each factor equal to zero and solve for the independent variable. You will need to use the quadratic formula if one of the factors is a nonfactorable quadratic.

For example, to solve the equation

$2x^4 + x^3 - 19x^2 - 14x + 24 = 0$, let $f(x) = 2x^4 + x^3 - 19x^2 - 14x + 24$. Possible values of x that make $f(x) = 0$ are numbers of the form $\frac{p}{q}$, where p is a factor of the constant term and q is a factor of the leading coefficient. Some possible values in this case are:

$$\frac{\pm 1, \pm 2, \pm 3, \pm 4, \pm 6, \pm 8, \pm 12, \pm 24}{\pm 1, \pm 2}$$

Since $f(-2) = 0$, by the factor theorem, $(x + 2)$ is a factor of $f(x)$. Determine $f(x) \div (x + 2)$ to find the other factor.

$$\begin{array}{r|rrrrr} -2 & 2 & 1 & -19 & -14 & 24 \\ & \downarrow & -4 & 6 & 26 & -24 \\ \hline & 2 & -3 & -13 & 12 & 0 \end{array}$$

$$\text{Now, } f(x) = (x + 2)(2x^3 - 3x^2 - 13x + 12).$$

Possible values of x that make the cubic polynomial 0 are numbers of the form:

$$\frac{\pm 1, \pm 2, \pm 3, \pm 4, \pm 6, \pm 12}{\pm 2}$$

Since $f(3) = 0$, $(x - 3)$ is also a factor of $f(x)$. Divide the cubic polynomial by $(x - 3)$ to determine the other factor.

$$\begin{array}{r|rrrr} 3 & 2 & -3 & -13 & 12 \\ & \downarrow & 6 & 9 & -12 \\ \hline & 2 & 3 & -4 & 0 \end{array}$$

$$\text{So, } f(x) = (x + 2)(x - 3)(2x^2 + 3x - 4).$$

$$x + 2 = 0 \text{ or } x - 3 = 0 \text{ or } 2x^2 + 3x - 4 = 0$$

$$x = -2 \text{ or } x = 3$$

Since $2x^2 + 3x - 4$ is not factorable, use the quadratic formula to determine the other zeros.

$$x = \frac{-3 \pm \sqrt{3^2 - 4(2)(-4)}}{2(2)}$$


$$x = \frac{-3 \pm \sqrt{41}}{4}$$

$$x \doteq -2.35 \text{ or } 0.85$$

The equation has four roots: $x = -2$, $x \doteq -2.35$, $x \doteq 0.85$, and $x = 3$.

A2: You can use a graphing strategy to find the zeros of a rearranged equation, or graph both sides of the equation separately and then determine the point(s) of intersection.

For example, to solve the equation $x^3 + 3x^2 - 7x + 4 = 3x^2 - 5x + 12$, enter both polynomials in the equation editor of the graphing calculator and then graph both corresponding functions. Use the intersect operation to determine the point of intersection of the two graphs.



The solution is $x \doteq 2.33$.

Q: How can you solve a linear inequality?

A: You solve a linear inequality using inverse operations in much the same way you would solve a linear equation. If at any time you multiply or divide the inequality by a negative number, you must reverse the inequality sign.

Study Aid

- See Lesson 4.2, Examples 1, 3, and 4.
- Try Mid-Chapter Review Questions 4 to 8.

PRACTICE Questions

Lesson 4.1


- Determine the solutions for each of the following.
 - $0 = -2x^3(2x - 5)(x - 4)^2$
 - $0 = (x^2 + 1)(2x + 4)(x + 2)$
 - $x^3 - 4x^2 = 7x - 10$
 - $0 = (x^2 - 2x - 24)(x^2 - 25)$
 - $0 = (x^3 + 2x^2)(x + 9)$
 - $-x^4 = -13x^2 + 36$
- Jude is diving from a cliff into the ocean. His height above sea level in metres is represented by the function $h(t) = -5(t - 0.3)^2 + 25$, where t is measured in seconds.
 - Expand the height function.
 - How high is the cliff?
 - When does Jude hit the water?
 - Determine where the function is negative. What is the significance of the negative values?
- Chris makes an open-topped box from a 30 cm by 30 cm piece of cardboard by cutting out equal squares from the corners and folding up the flaps to make the sides. What are the dimensions of each square, to the nearest hundredth of a centimetre, so that the volume of the resulting box is 1000 cm^3 ?

Lesson 4.2

- Solve the following inequalities algebraically and plot the solution on a number line.
 - $2x - 4 < 3x + 7$
 - $-x - 4 \leq x + 4$
 - $-2(x - 4) \geq 16$
 - $2(3x - 7) > 3(7x - 3)$
- Solve and state your solution using interval notation.

$$2x < \frac{3x + 6}{2} \leq 4 + 2x$$

- Create a linear inequality with both a constant and a linear term on each side and that has each of the following as a solution.
 - $x > 7$
 - $x \in (-\infty, -8)$
 - $-1 \leq x \leq 7$
 - $x \in [3, \infty)$
- Consider the following functions.



- Find the equations of the lines depicted.
 - Solve the inequality $f(x) < g(x)$ by examining the graph.
 - Confirm your solution by solving the inequality algebraically.
- The New Network cell phone company charges \$20 a month for service and \$0.02 per minute of talking time. The My Mobile company charges \$15 a month for service and \$0.03 per minute of talking time.
 - Write expressions for the total bill of each company.
 - Set up an inequality that can be used to determine for what amount of time (in minutes) My Mobile is the better plan.
 - Solve your inequality.
 - Why did you have to put a restriction on the algebraic solution from part c)?

4.3

Solving Polynomial Inequalities

GOAL

Solve polynomial inequalities.

YOU WILL NEED

- graphing calculator or graphing software

LEARN ABOUT the Math



The elevation of a hiking trail is modelled by the function $h(x) = 2x^3 + 3x^2 - 17x + 12$, where h is the height measured in metres above sea level and x is the horizontal position from a ranger station measured in kilometres. If x is negative, the position is to the west of the station, and if x is positive, the position is to the east. Since the trail extends 4.2 km to the west of the ranger station and 4 km to the east, the model is accurate where $x \in [-4.2, 4]$.

? How can you determine which sections of the trail are above sea level?

EXAMPLE 1

Selecting a strategy to solve the problem

At what distances from the ranger station is the trail above sea level?

Solution A: Using an algebraic strategy and a number line

$$2x^3 + 3x^2 - 17x + 12 > 0$$

The trail is above sea level when the height is positive, i.e., $h(x) > 0$. Write the mathematical model using a **polynomial inequality**.

polynomial inequality

an inequality that contains a polynomial expression (e.g., $5x^3 + 3x^2 - 6x \leq 2$)

$$h(1) = 2(1)^3 + 3(1)^2 - 17(1) + 12$$

$$= 0 \text{ so } (x - 1) \text{ is a factor of } h(x)$$

$$\begin{array}{r|rrrr} 1 & 2 & 3 & -17 & 12 \\ & \downarrow & & & \\ & 2 & 5 & -12 & 0 \end{array}$$

$$h(x) = (x - 1)(2x^2 + 5x - 12)$$

$$0 = (x - 1)(2x - 3)(x + 4)$$


$$x = 1, x = \frac{3}{2}, \text{ or } x = -4$$

Factor the corresponding function $y = h(x)$ to locate the x -intercepts. Use the factor theorem to determine the first factor.

Set $h(x) = 0$. Set each factor equal to 0 and solve.

The x -intercepts are at -4 , 1 , and $\frac{3}{2}$. These numbers divide the domain of real numbers into four intervals:

$$x < -4, -4 < x < 1, 1 < x < \frac{3}{2}, x > \frac{3}{2}$$



Draw a number line and test points in each interval to see whether the function has a positive or negative value.

Interval	$x < -4$	$-4 < x < 1$	$1 < x < \frac{3}{2}$	$x > \frac{3}{2}$
Value of $h(x)$	$h(-5) = -78$	$h(0) = 12$	$h(1.2) \doteq -0.6$	$h(2) = 6$
Is $h(x) > 0$?	no	yes	no	yes

$$h(x) > 0 \text{ when } -4 < x < 1 \text{ and } x > \frac{3}{2}$$

Identify the intervals where $h(x)$ is positive.

The hiking trail is above sea level from 4 km west of the ranger station to 1 km east, and for distances more than 1.5 km east.

Write a concluding statement.

Solution B: Using a graphing strategy

$$2x^3 + 3x^2 - 17x + 12 > 0$$

The trail is above sea level when the height is positive, i.e., $h(x) > 0$.

$$h(1) = 2(1)^3 + 3(1)^2 - 17(1) + 12$$

$$= 0 \text{ so } (x - 1) \text{ is a factor of } h(x)$$

$$\begin{array}{r|rrrr} 1 & 2 & 3 & -17 & 12 \\ & \downarrow & & & \\ & 2 & 5 & -12 & 0 \end{array}$$

Factor the corresponding function $y = h(x)$ to locate the x -intercepts. Use the factor theorem to determine the first factor.


$$h(x) = (x - 1)(2x^2 + 5x - 12)$$

$$0 = (x - 1)(2x - 3)(x + 4)$$

$$x = 1, x = \frac{3}{2}, \text{ or } x = -4$$

The x -intercepts are at -4 , 1 , and $\frac{3}{2}$.

$$h(0) = 12$$



The graph lies above the x -axis on the intervals $-4 < x < 1$ and $x > \frac{3}{2}$.

The hiking trail is above sea level from 4 km west of the ranger station to 1 km east, and for distances beyond 1.5 km to the east of the ranger station.

Set $h(x) = 0$.
Set each factor equal to 0 and solve.

The y -intercept occurs when $x = 0$.

Analyze the function and draw a sketch of $h(x)$. Plot the x - and y -intercepts. Because the leading coefficient of the function is positive and the degree of the function is odd, the graph has opposite end behaviours. The graph must start in the third quadrant and proceed to the first quadrant. Estimate the location of the turning points.

Determine the intervals where $h(x) > 0$.

Write a concluding statement that answers the question.

Reflecting

- A. When solving a polynomial inequality, which steps are the same as those used when solving a polynomial equation?
- B. What additional steps must be taken when solving a polynomial inequality?
- C. The zeros of $y = h(x)$ were used to identify the intervals where $h(x)$ was positive and negative but were not included in the solution set of $h(x) > 0$. Explain why.
- D. How could you verify the solution set to the polynomial inequality using graphing technology?

APPLY the Math

EXAMPLE 2

Selecting tools and strategies to solve a factorable polynomial inequality

Solve the inequality $x^3 - 2x^2 + 5x + 20 \geq 2x^2 + 14x - 16$.

Solution A: Using algebra and a factor table

$$x^3 - 2x^2 + 5x + 20 \geq 2x^2 + 14x - 16$$

$$x^3 - 4x^2 - 9x + 36 \geq 0$$

Use inverse operations to make the right side of the inequality equal to zero.

$$x^2(x - 4) - 9(x - 4) \geq 0$$

$$(x - 4)(x^2 - 9) \geq 0$$

Factor the polynomial on the left by grouping.

$$(x - 4)(x - 3)(x + 3) \geq 0$$

$$(x - 4)(x - 3)(x + 3) = 0$$

Determine the roots of the corresponding polynomial equation.

The roots are -3 , 3 , and 4 . These numbers divide the real numbers into four intervals:

$$x < -3, -3 < x < 3, 3 < x < 4, x > 4$$



	$x < -3$	$-3 < x < 3$	$3 < x < 4$	$x > 4$
$(x - 4)$	—	—	—	+
$(x - 3)$	—	—	+	+
$(x + 3)$	—	+	+	+
their product	$(-)(-)(-) = -$	$(-)(-)(+) = +$	$(-)(+)(+) = -$	$(+)(+)(+) = +$

Create a table to consider the sign of each factor in each of the intervals and examine the sign of their product. In this case, the intervals that correspond to a positive product are the solutions to the polynomial inequality.



$$x^3 - 2x^2 + 5x + 20 \geq 2x^2 + 14x - 16 \text{ when } -3 \leq x \leq 3 \text{ or } x \geq 4.$$

Write a concluding statement.



Solution B: Using graphing technology



Graph each side of the inequality as a separate function. Bold the graph of the second function (the quadratic) so you can distinguish one from the other. Experiment with different window settings to make the intersecting parts of the graph visible.



The two graphs intersect somewhere to the left of the y -axis and intersect twice to the right of the y -axis. Use the intersect operation to determine all points of intersection.



You can see on the graph that the cubic function lies above the quadratic function in the interval $-3 \leq x \leq 3$ or $x \geq 4$.

The two functions intersect at $(-3, -40)$, $(3, 44)$, and $(4, 72)$.

$$x^3 - 10x^2 + 15x + 11 \geq -8x + 26$$


when $x \in [-3, 3]$ or $x \in [4, \infty)$.

Write a concluding statement.

EXAMPLE 3

Selecting a strategy to solve a polynomial inequality that is unfactorable

The height of one section of the roller coaster can be described by the polynomial function $h(x) = \frac{1}{4\,000\,000}x^2(x - 30)^2(x - 55)^2$, where h is the height, measured in metres, and x is the position from the start, measured in metres along the ground.



When will the roller coaster car be more than 9 m above the ground?





Solution



Solve

$$\frac{1}{4\,000\,000}x^2(x-30)^2(x-55)^2 > 9$$



In this case, the solution set corresponds to all values of x where $h(x) > 9$. Using an algebraic approach involving factoring would be tedious, so use a graphing strategy.



Graph the function $h(x) = \frac{1}{4\,000\,000}x^2(x-30)^2(x-55)^2$ and the line $y = 9$ on the graphing calculator and locate intervals where the roller coaster is higher than 9 m. On the graph, this will correspond to when $Y1 > Y2$.



Determine the four points of intersection of the height function and the horizontal line.



The four points where the height function and the horizontal line intersect are approximately $(4.7, 9)$, $(21.7, 9)$, $(40, 9)$, and $(48.1, 9)$.

The roller coaster will be more than 9 m above the ground when it is between 4.7 m and 21.7 m from the starting point and between 40 m and 48.1 m from the starting point, as measured along the ground.

$Y1 > Y2$ when $4.7 < x < 21.7$ or $40 < x < 48.1$.

In Summary

Key Idea



- To solve a polynomial inequality algebraically, you must first determine the roots of the corresponding polynomial equation. Then you must consider the sign of the polynomial in each of the intervals created by these roots. The solution set is determined by the interval(s) that satisfy the given inequality.



Need to Know

- Some polynomial inequalities can be solved algebraically by
 - using inverse operations to move all terms to one side of the inequality
 - factoring the polynomial to determine the zeros of the corresponding polynomial equation
 - using a number line, a graph, or a factor table to determine the intervals on which the polynomial is positive or negative
- All polynomial inequalities can be solved using graphing technology by
 - graphing each side of the inequality as a separate function
 - determining the intersection point(s) of the functions
 - examining the graph to determine the intervals where one function is above or below the other, as required
 or
 - creating an equivalent inequality with zero on one side
 - identifying the intervals created by the zeros of the graph of the new function
 - finding where the graph lies above the x -axis (where $f(x) > 0$) or below (where $f(x) < 0$), as required

CHECK Your Understanding

- Solve each of the following using a number line strategy. Express your answers using set notation.
 - $(x + 2)(x - 3)(x + 1) \geq 0$
 - $-2(x - 2)(x - 4)(x + 3) < 0$
 - $(x - 3)(5x + 2)(4x - 3) < 0$
 - $(x - 5)(4x + 1)(2x - 5) \geq 0$
- For each graph shown, determine where $f(x) \leq 0$. Express your answers using interval notation.









3. If $f(x) = 2x^3 - x^2 + 3x + 10$ and $g(x) = x^3 + 3x^2 + 2x + 4$, determine when $f(x) > g(x)$ using a factor table strategy.
4. Solve the inequality $x^3 - 7x^2 + 4x + 12 > x^2 - 4x - 9$ using a graphing calculator.

PRACTISING

5. For each of the following polynomial functions, state the intervals where $f(x) > 0$.

K




6. Solve the following inequalities.
- $(x - 1)(x + 1) > 0$
 - $(x + 3)(x - 4) < 0$
 - $(2x + 1)(x - 5) \geq 0$
 - $-3x(x + 7)(x - 2) < 0$
 - $(x - 3)(x + 1) + (x - 3)(x + 2) \geq 0$
 - $2x(x + 4) - 3(x + 4) \leq 0$

7. Solve the following inequalities algebraically. Confirm your answer with a graph.


- $x^2 - 6x + 9 \geq 16$
- $x^4 - 8x < 0$
- $x^3 + 4x^2 + x \leq 6$
- $x^4 - 5x^2 + 4 > 0$
- $3x^3 - 3x^2 - 2x \leq 2x^3 - x^2 + x$
- $x^3 - x^2 - 3x + 3 > -x^3 + 2x + 5$

8. For the following pair of functions, determine when $f(x) < g(x)$.



- Consider $x^3 + 11x^2 + 18x + 10 > 10$.
 - What is the equation of the corresponding function that could be graphed and used to solve this inequality?
 - Explain how the graph of the corresponding function can be used in this case to solve the inequality.
 - Solve this inequality algebraically.
- Determine an expression for $f(x)$ in which $f(x)$ is a quartic function, $f(x) > 0$ when $-2 < x < 1$, $f(x) \leq 0$ when $x < -2$ or $x > 1$, $f(x)$ has a double root when $x = 3$, and $f(-1) = 96$.
- The viscosity, v , of oil used in cars is related to its temperature, t , by the formula $v = -t^3 - 6t^2 + 12t + 50$, where each unit of t is equivalent to 50°C .
 - Graph the function on your graphing calculator.
 - Determine the temperature range for which $v > 0$ to two decimal places.
 - Determine the temperature ranges for which $15 < v < 20$ to two decimal places.

12. A rock is tossed from a platform and follows a parabolic path through the air. The height of the rock in metres is given by $h(t) = -5t^2 + 12t + 14$, where t is measured in seconds.
- How high is the rock off the ground when it is thrown?
 - How long is the rock in the air?
 - For what times is the height of the rock greater than 17 m?
 - How long is the rock above a height of 17 m?
13. An open-topped box can be made from a sheet of aluminium measuring 50 cm by 30 cm by cutting congruent squares from the four corners and folding up the sides. Write a polynomial function to represent the volume of such a box. Determine the range of side lengths that are possible for each square that is cut out and removed that result in a volume greater than 4000 cm^3 .



14. a) Without a calculator, explain why the inequality $2x^{24} + x^4 + 15x^2 + 80 < 0$ has no solution.
 b) Without a calculator, explain why $-4x^{12} - 7x^6 + 9x^2 + 20 < 30 + 11x^2$ has a solution of $-\infty < x < \infty$.
15. Explain why the following solution strategy fails, and then solve the inequality correctly.
 Solve: $(x + 1)(x - 2) > (x + 1)(-x + 6)$.
 Divide both sides by $x + 1$ and get $x - 2 > -x + 6$.
 Add x to both sides: $2x - 2 > 6$.
 Add 2 to both sides: $2x > 8$.
 Divide both sides by 2: $x > 4$.
16. Create a concept web that illustrates all of the different methods you could use to solve a polynomial inequality.

Extending

17. Use what you know about the factoring method to solve the following inequalities.
- $\frac{x^2 + x - 12}{x^2 + 5x + 6} < 0$
 - $\frac{x^2 - 25}{x^3 + 6x^2 + 5x} > 0$
18. Solve the inequality $(x + 1)(x - 2)(2^x) \geq 0$ algebraically.





Chapter

5

Rational Functions, Equations, and Inequalities

► GOALS

You will be able to

- Graph the reciprocal functions of linear and quadratic functions
- Identify the key characteristics of rational functions from their equations and use these characteristics to sketch their graphs
- Solve rational equations and inequalities with and without graphing technology
- Determine average and instantaneous rates of change in situations that are modelled by rational functions

? When polluted water flows into a clean pond, how does the concentration of pollutant in the pond change over time? What type of function would model this change?

Study Aid

- For help, see the Review of Essential Skills found at the Nelson Advanced Functions website.

Question	Appendix
1	R-3
2, 3, 4, 8	R-4
7	R-8

SKILLS AND CONCEPTS You Need

- Factor each expression.
 - $x^2 - 3x - 10$
 - $3x^2 + 12x - 15$
 - $16x^2 - 49$
 - $9x^2 - 12x + 4$
 - $3a^2 + a - 30$
 - $6x^2 - 5xy - 21y^2$
- Simplify each expression. State any restrictions on the variables, if necessary.
 - $\frac{12 - 8s}{4}$
 - $\frac{6m^2n^4}{18m^3n}$
 - $\frac{9x^3 - 12x^2 - 3x}{3x}$
 - $\frac{25x - 10}{5(5x - 2)^2}$
 - $\frac{x^2 + 3x - 18}{9 - x^2}$
 - $\frac{a^2 + 4ab - 5b^2}{2a^2 + 7ab - 15b^2}$
- Simplify each expression, and state any restrictions on the variable.
 - $\frac{3}{5} \times \frac{7}{9}$
 - $\frac{2x}{5} \div \frac{x^2}{15}$
 - $\frac{x^2 - 4}{x - 3} \div \frac{x + 2}{12 - 4x}$
 - $\frac{x^3 + 4x^2}{x^2 - 1} \times \frac{x^2 - 5x + 6}{x^2 - 3x}$
- Simplify each expression, and state any restrictions on the variable.
 - $\frac{2}{3} + \frac{6}{7}$
 - $\frac{3x}{4} + \frac{5x}{6}$
 - $\frac{1}{x} + \frac{4}{x^2}$
 - $\frac{5}{x - 3} - \frac{2}{x}$
 - $\frac{2}{x - 5} + \frac{y}{x^2 - 25}$
 - $\frac{6}{a^2 - 9a + 20} - \frac{8}{a^2 - 2a - 15}$
- Solve and check.
 - $\frac{5x}{8} = \frac{15}{4}$
 - $\frac{x}{4} + \frac{1}{3} = \frac{5}{6}$
 - $\frac{4x}{5} - \frac{3x}{10} = \frac{3}{2}$
 - $\frac{x + 1}{2} - \frac{2x - 1}{3} = -1$
- Sketch the graph of the reciprocal function $f(x) = \frac{1}{x}$ and describe its characteristics. Include the domain and range, as well as the equations of the asymptotes.

7. List the transformations that need to be applied to $y = \frac{1}{x}$ to graph each of the following reciprocal functions. Then sketch the graph.
- a) $f(x) = \frac{1}{x+3}$ c) $f(x) = -\frac{1}{2x} - 3$
- b) $f(x) = \frac{2}{x-1}$ d) $f(x) = \frac{2}{-3(x-2)} + 1$
8. Describe the steps that are required to divide two rational expressions. Use your description to simplify $\frac{9y^2 - 4}{4y - 12} \div \frac{9y^2 + 12y + 4}{18 - 6y}$.

APPLYING What You Know

Painting Houses

Tony can paint the exterior of a house in six working days. Rebecca takes nine days to complete the same painting job.

- ?** How long will Rebecca and Tony take to paint a similar house, if they work together?
- A. What fraction of the job can Tony complete in one day? What fraction of the job can Rebecca complete?
- B. Write a numerical expression to represent the fraction of the job that Rebecca and Tony can complete in one day, if they work together.
- C. Let x represent the number of days that Rebecca and Tony, working together, will take to complete the job. Explain why $\frac{1}{x}$ represents the fraction of the job Rebecca and Tony will complete in one day when they work together.
- D. Use your answers for parts B and C to write an equation. Determine the **lowest common denominator** for the **rational expressions** in your equation. Rewrite the equation using the lowest common denominator.
- E. Solve the equation you wrote in part D by collecting like terms and comparing the numerators on the two sides of the equation.
- F. What is the amount of time Rebecca and Tony will take to paint a similar house, when they work together?




5.1

Graphs of Reciprocal Functions

YOU WILL NEED

- graph paper
- coloured pencils or pens
- graphing calculator or graphing software



GOAL

Sketch the graphs of reciprocals of linear and quadratic functions.

INVESTIGATE the Math

Owen has noted some connections between the graphs of $f(x) = x$ and its reciprocal function $g(x) = \frac{1}{x}$.

- Both graphs are in the same quadrants for the same x -values.
- When $f(x) = 0$, there is a vertical asymptote for $g(x)$.
- $f(x)$ is always increasing, and $g(x)$ is always decreasing.

? How are the graphs of a function and its reciprocal function related?

- Explain why the graphs of $f(x) = x$ and $g(x) = \frac{1}{x}$ are in the same quadrants over the same intervals. Does this relationship hold for $m(x) = -x$ and $n(x) = -\frac{1}{x}$? Does this relationship hold for any function and its reciprocal function? Explain.
- What graphical characteristic in the reciprocal function do the zeros of the original function correspond to? Explain.
- Explain why the reciprocal function $g(x) = \frac{1}{x}$ is decreasing when $f(x) = x$ is increasing. Does this relationship hold for $n(x) = -\frac{1}{x}$ and $m(x) = -x$? Explain how the increasing and decreasing intervals of a function and its reciprocal are related.
- What are the y -coordinates of the points where $f(x)$ and $g(x)$ intersect? Will the points of intersection for any function and its reciprocal always have the same y -coordinates? Explain.
- Explain why the graph of $g(x)$ has a horizontal asymptote. What is the equation of this asymptote? Will all reciprocal functions have the same horizontal asymptote? Explain.


- F. On graph paper, draw the graph of $p(x) = x^2 - 4$. In a table like the one below, note the characteristics of the graph of $p(x)$ and use this information to help you determine the characteristics of the reciprocal function $q(x) = \frac{1}{x^2 - 4}$.

Characteristics	$p(x) = x^2 - 4$	$q(x) = \frac{1}{x^2 - 4}$
zeros and/or vertical asymptotes		
interval(s) on which the graph is above the x-axis (all values of the function are positive)		
interval(s) on which the graph is below the x-axis (all values of the function are negative)		
interval(s) on which the function is increasing		
interval(s) on which the function is decreasing		
point(s) where the y-value is 1		
point(s) where the y-value is -1		

- G. On the same graph, draw the vertical asymptotes for the reciprocal function. Then use the rest of the information determined in part F to draw the graph for $q(x) = \frac{1}{x^2 - 4}$.
- H. Verify your graphs by entering $p(x)$ and $q(x)$ in a graphing calculator using the “friendly” window setting shown.
- I. Repeat parts F to H for the following pairs of functions.
- $p(x) = x + 2$ and $q(x) = \frac{1}{x + 2}$
 - $p(x) = 2x - 3$ and $q(x) = \frac{1}{2x - 3}$
 - $p(x) = (x - 2)(x + 3)$ and $q(x) = \frac{1}{(x - 2)(x + 3)}$
 - $p(x) = (x - 1)^2$ and $q(x) = \frac{1}{(x - 1)^2}$
- J. Write a summary of the relationships between the characteristics of the graphs of
- a linear function and its reciprocal function
 - a quadratic function and its reciprocal function

Tech Support

On a graphing calculator, the length of the display screen contains 94 pixels, and the width contains 62 pixels. When the domain, $X_{\max} - X_{\min}$, is cleanly divisible by 94, and the range, $Y_{\max} - Y_{\min}$, is cleanly divisible by 62, the window is friendly. This means that you can trace without using “ugly” decimals. A friendly window is useful when working with rational functions.



Use brackets when entering reciprocal functions in the $Y =$ editor of a graphing calculator. For example, to graph the function $f(x) = \frac{1}{x^2 - 4}$, enter $Y1 = \frac{1}{(x^2 - 4)}$.

Reflecting

- K. How did knowing the positive/negative intervals and the increasing/decreasing intervals for $p(x) = x^2 - 4$ help you draw the graph for $p(x) = \frac{1}{x^2 - 4}$?
- L. Why are some numbers in the domain of a function excluded from the domain of its reciprocal function? What graphical characteristic of the reciprocal function occurs at these values?
- M. What common characteristics are shared by all reciprocals of linear and quadratic functions?

APPLY the Math

EXAMPLE 1

Connecting the characteristics of a linear function to its corresponding reciprocal function

Given the function $f(x) = 2 - x$,

- a) determine the domain and range, intercepts, positive/negative intervals, and increasing/decreasing intervals
- b) use your answers for part a) to sketch the graph of the reciprocal function

Solution

- a) $f(x) = 2 - x$ is a linear function.

$$D = \{x \in \mathbf{R}\}$$


$$R = \{y \in \mathbf{R}\}$$

From the equation, the y -intercept is 2.

$$f(x) = 0 \text{ when } 0 = 2 - x$$

$$x = 2$$

The x -intercept is 2.



$f(x)$ is positive when $x \in (-\infty, 2)$ and negative when $x \in (2, \infty)$.

$f(x)$ is decreasing when $x \in (-\infty, \infty)$.

The domain and range of most linear functions are the set of real numbers.

A linear function $f(x) = mx + b$ has y -intercept b .

The x -intercept occurs where $f(x) = 0$.

Sketch the graph of $f(x)$ to determine the positive and negative intervals. The line $y = 2 - x$ is above the x -axis for all x -values less than 2 and below the x -axis for all x -values greater than 2.

This is a linear function with a negative slope, so it is decreasing over its entire domain.

b) The reciprocal function is $g(x) = \frac{1}{2-x}$.

$$D = \{x \in \mathbf{R} \mid x \neq 2\}$$

$$R = \{y \in \mathbf{R} \mid y \neq 0\}$$

The y -intercept is 0.5.

The vertical asymptote is $x = 2$
and the horizontal asymptote is $y = 0$.

The reciprocal function is positive
when $x \in (-\infty, 2)$ and negative
when $x \in (2, \infty)$.

It is increasing when $x \in (-\infty, 2)$
and when $x \in (2, \infty)$.

The graph of $g(x) = \frac{1}{2-x}$ intersects
the graph of $g(x) = 2-x$ at $(1, 1)$
and $(3, -1)$.


All the y -values of points on the reciprocal function are reciprocals of the y -values on the original function.

There is a vertical asymptote at the zero of the original function.
The reciprocals of a linear function always have the x -axis as a horizontal asymptote.

The positive/negative intervals are always the same for both functions.

Because the original function is always decreasing, the reciprocal function is always increasing.

The reciprocal of 1 is 1, and the reciprocal of -1 is -1 . Thus, the two graphs intersect at any points with these y -values.



Use all this information to sketch the graph of the reciprocal function.

EXAMPLE 2

Connecting the characteristics of a quadratic function to its corresponding reciprocal function

Given the function $f(x) = 9 - x^2$

- determine the domain and range, intercepts, positive/negative intervals, and increasing/decreasing intervals
- use your answers for part a) to sketch the graph of the reciprocal function

Solution

a) $f(x) = 9 - x^2$ is a quadratic function.

$$D = \{x \in \mathbf{R}\}$$

The domain of a quadratic function is the set of real numbers.

$f(0) = 9$, so the y -intercept is 9.

$$R = \{y \in \mathbf{R} \mid y \leq 9\}$$

The graph of $f(x)$ is a parabola that opens down. The vertex is at $(0, 9)$, so $y \leq 9$.


$$f(x) = 0 \Rightarrow 9 - x^2 = 0$$

$$(3 - x)(3 + x) = 0$$

$$x = \pm 3$$

Factor and determine the x -intercepts.

The x -intercepts are -3 and 3 .



The parabola $y = 9 - x^2$ is above the x -axis for x -values between -3 and 3 . The graph is below the x -axis for x -values less than -3 and for x -values greater than 3 .

$f(x)$ is positive when $x \in (-3, 3)$ and negative when $x \in (-\infty, -3)$ and when $x \in (3, \infty)$.

$f(x)$ is increasing when $x \in (-\infty, 0)$ and decreasing when $x \in (0, \infty)$.

The y -values increase as x increases from $-\infty$ to 0 . The y -values decrease as x increases from 0 to ∞ .

b) The reciprocal function is $g(x) = \frac{1}{9 - x^2}$.

The vertical asymptotes are $x = -3$ and $x = 3$.

$$D = \{x \in \mathbf{R} \mid x \neq \pm 3\}$$

Vertical asymptotes occur at each zero of the original function, so these numbers must be excluded from the domain.

The horizontal asymptote is $y = 0$.

The y -intercept is $\frac{1}{9}$.

The reciprocals of all quadratic functions have the x -axis as a horizontal asymptote. The y -intercept of the original function is 9 , so the y -intercept of the reciprocal function is $\frac{1}{9}$.

There is a local minimum value at $\left(0, \frac{1}{9}\right)$.

$$R = \left\{y \in \mathbf{R} \mid y < 0 \text{ or } y \geq \frac{1}{9}\right\}$$

When the original function has a local maximum point, the reciprocal function has a corresponding local minimum point.




The reciprocal function is positive when $x \in (-3, 3)$ and negative when $x \in (-\infty, -3)$ and when $x \in (3, \infty)$. It is decreasing when $x \in (-\infty, -3)$ and when $x \in (-3, 0)$, and increasing when $x \in (0, 3)$ and when $x \in (3, \infty)$.

The positive/negative intervals are always the same for both functions. Where the original function is decreasing, excluding the zeros, the reciprocal function is increasing (and vice versa).

$$\begin{aligned} f(x) = 1 \text{ when } 9 - x^2 = 1 & \quad \text{and} \quad f(x) = -1 \text{ when } 9 - x^2 = -1 \\ -x^2 = 1 - 9 & \quad -x^2 = -1 - 9 \\ -x^2 = -8 & \quad -x^2 = -10 \\ x^2 = 8 & \quad x^2 = 10 \\ x = \pm 2\sqrt{2} & \quad x = \pm \sqrt{10} \end{aligned}$$

A function and its reciprocal intersect at points where $y = \pm 1$. Solve the corresponding equations to determine the x-coordinates of the points of intersection.

The graph of $g(x) = \frac{1}{9 - x^2}$ intersects the graph of $f(x) = 9 - x^2$ at $(-2\sqrt{2}, 1)$, $(2\sqrt{2}, 1)$ and at $(-\sqrt{10}, -1)$, $(\sqrt{10}, -1)$.



Use all this information to sketch the graph of the reciprocal function.

In Summary

Key Idea

- You can use key characteristics of the graph of a linear or quadratic function to graph the related reciprocal function.


Need to Know

- All the y-coordinates of a reciprocal function are the reciprocals of the y-coordinates of the original function.
- The graph of a reciprocal function has a vertical asymptote at each zero of the original function.
- A reciprocal function will always have $y = 0$ as a horizontal asymptote if the original function is linear or quadratic.
- A reciprocal function has the same positive/negative intervals as the original function.

(continued)


- Intervals of increase on the original function are intervals of decrease on the reciprocal function. Intervals of decrease on the original function are intervals of increase on the reciprocal function.
- If the range of the original function includes 1 and/or -1 , the reciprocal function will intersect the original function at a point (or points) where the y -coordinate is 1 or -1 .
- If the original function has a local minimum point, the reciprocal function will have a local maximum point at the same x -value (and vice versa).

A linear function and its reciprocal



Both functions are negative when $x \in (-\infty, -1)$ and positive when $x \in (-1, \infty)$. The original function is increasing when $x \in (-\infty, \infty)$. The reciprocal function is decreasing when $x \in (-\infty, -1)$ or $(-1, \infty)$.


A quadratic function and its reciprocal



Both functions are negative when $x \in (-3, 1)$ and positive when $x \in (-\infty, -3)$ or $(1, \infty)$. The original function is decreasing when $x \in (-\infty, -1)$ and increasing when $x \in (-1, \infty)$. The reciprocal function is increasing when $x \in (-\infty, -3)$ or $(-3, -1)$ and decreasing when $x \in (-1, 1)$ or $(1, \infty)$.

CHECK Your Understanding

1. Match each function with its equation on the next page. Then identify which function pairs are reciprocals.



$$\text{A } y = \frac{1}{-(x-2)^2 - 1}$$

$$\text{D } y = x^2 - 1$$

$$\text{B } y = \frac{1}{x^2 - 1}$$

$$\text{E } y = -(x-2)^2 - 1$$

$$\text{C } y = \frac{1}{2x - 5}$$

$$\text{F } y = 2x - 5$$

2. For each pair of functions, determine where the zeros of the original function occur and state the equations of the vertical asymptotes of the reciprocal function, if possible.

a) $f(x) = x - 6, g(x) = \frac{1}{x - 6}$

b) $f(x) = 3x + 4, g(x) = \frac{1}{3x + 4}$

c) $f(x) = x^2 - 2x - 15, g(x) = \frac{1}{x^2 - 2x - 15}$

d) $f(x) = 4x^2 - 25, g(x) = \frac{1}{4x^2 - 25}$

e) $f(x) = x^2 + 4, g(x) = \frac{1}{x^2 + 4}$

f) $f(x) = 2x^2 + 5x + 3, g(x) = \frac{1}{2x^2 + 5x + 3}$

3. Sketch the graph of each function. Use your graph to help you sketch the graph of the reciprocal function.

a) $f(x) = 5 - x$ b) $f(x) = x^2 - 6x$

PRACTISING

4. a) Copy and complete the following table.

x	-4	-3	-2	-1	0	1	2	3	4	5	6	7
$f(x)$	16	14	12	10	8	6	4	2	0	-2	-4	-6
$\frac{1}{f(x)}$												

b) Sketch the graphs of $y = f(x)$ and $y = \frac{1}{f(x)}$.

c) Find equations for $y = f(x)$ and $y = \frac{1}{f(x)}$.

5. State the equation of the reciprocal of each function, and determine the equations of the vertical asymptotes of the reciprocal. Verify your results using graphing technology.





a) $f(x) = 2x$ e) $f(x) = -3x + 6$

b) $f(x) = x + 5$ f) $f(x) = (x - 3)^2$

c) $f(x) = x - 4$ g) $f(x) = x^2 - 3x - 10$

d) $f(x) = 2x + 5$ h) $f(x) = 3x^2 - 4x - 4$

6. Sketch the graph of the reciprocal of each function.



7. Sketch each pair of graphs on the same axes. State the domain and range of each reciprocal function.

a) $y = 2x - 5, y = \frac{1}{2x - 5}$

b) $y = 3x + 4, y = \frac{1}{3x + 4}$

8. Draw the graph of $y = f(x)$ and $y = \frac{1}{f(x)}$ on the same axes.

a) $f(x) = x^2 - 4$

d) $f(x) = (x + 3)^2$

b) $f(x) = (x - 2)^2 - 3$

e) $f(x) = x^2 + 2$

c) $f(x) = x^2 - 3x + 2$

f) $f(x) = -(x + 4)^2 + 1$

9. For each function, determine the domain and range, intercepts, positive/negative intervals, and increasing/decreasing intervals. State the equation of the reciprocal function. Then sketch the graphs of the original and reciprocal functions on the same axes.

a) $f(x) = 2x + 8$


c) $f(x) = x^2 - x - 12$

b) $f(x) = -4x - 3$

d) $f(x) = -2x^2 + 10x - 12$

10. Why do the graphs of reciprocals of linear functions always have vertical asymptotes, but the graphs of reciprocals of quadratic functions sometimes do not? Provide sketches of three different reciprocal functions to illustrate your answer.

11. An equation of the form $y = \frac{k}{x^2 + bx + c}$ has a graph that closely matches the graph shown. Find the equation. Check your answer using graphing technology.



12. A chemical company is testing the effectiveness of a new cleaning solution for killing bacteria. The test involves introducing the solution into a sample that contains approximately 10 000 bacteria. The number of bacteria remaining, $b(t)$, over time, t , in seconds is given by the equation $b(t) = 10\,000 \frac{1}{t}$.

- How many bacteria will be left after 20 s?
 - After how many seconds will only 5000 bacteria be left?
 - After how many seconds will only one bacterium be left?
 - This model is not always accurate. Determine what sort of inaccuracies this model might have. Assume that the solution was introduced at $t = 0$.
 - Based on these inaccuracies, what should the domain and range of the equation be?
13. Use your graphing calculator to explore and then describe the key characteristics of the family of reciprocal functions of the form $g(x) = \frac{1}{x + n}$. Make sure that you include graphs to support your descriptions.
- State the domain and range of $g(x)$.
 - For the family of functions $f(x) = x + n$, the y -intercept changes as the value of n changes. Describe how the y -intercept changes and how this affects $g(x)$.
 - If graphed, at what point would the two graphs $f(x)$ and $g(x)$ intersect?
14. Due to a basketball tournament, your friend has missed this class.

- C** Write a concise explanation of the steps needed to graph a reciprocal function using the graph of the original function (without using graphing technology). Use an example, and explain the reason for each step.

Extending

15. Sketch the graphs of the following reciprocal functions.


a) $y = \frac{1}{\sqrt{x}}$

c) $y = \frac{1}{2^x}$

b) $y = \frac{1}{x^3}$

d) $y = \frac{1}{\sin x}$

16. Determine the equation of the function in the graph shown.



5.2

Exploring Quotients of Polynomial Functions

YOU WILL NEED

- graph paper
- coloured pencils or pens
- graphing calculator or graphing software



GOAL

Explore graphs that are created by dividing polynomial functions.



EXPLORE the Math

Each row shows the graphs of two polynomial functions.



A.



B.




C.



rational function

a function that can be expressed as $f(x) = \frac{p(x)}{q(x)}$, where $p(x)$ and $q(x)$ are polynomial functions, $q(x) \neq 0$ (e.g., $f(x) = \frac{3x^2 - 1}{x + 1}$, $x \neq -1$, and $f(x) = \frac{1 - x}{x^2}$, $x \neq 0$, are rational functions, but $f(x) = \frac{1 + x}{\sqrt{2 - x}}$, $x \neq 2$, is not because its denominator is not a polynomial)



? What are the characteristics of the graphs that are created by dividing two polynomial functions?

- A. Using the given functions, write the equation of the rational function $y = \frac{f(x)}{g(x)}$. Enter this equation into Y1 of the equation editor of a graphing calculator. Graph this equation using the window settings shown, and draw a sketch.


- B.** Describe the characteristics of the graph you created in part A by answering the following questions:
- i) Where are the zeros?
 - ii) Are there any asymptotes? If so, where are they?
 - iii) What are the domain and range of this function?
 - iv) Is it a **continuous function**? Explain.
 - v) Are there any values of $y = \frac{f(x)}{g(x)}$ that are undefined? What feature(s) of the graph is (are) related to these values?
 - vi) Describe the end behaviours of this function.
 - vii) Is the resulting graph a function? Explain.
- C.** Write the equation defined by $y = \frac{g(x)}{f(x)}$. Predict how the graph of this function will differ from the graph of $y = \frac{f(x)}{g(x)}$. Graph this function using your graphing calculator, and draw a sketch.
- D.** Describe the characteristics of the graph you created in part C by answering the questions in part B.
- E.** Repeat parts A through D for the functions in the other two rows.
- F.** Using graphing technology, and the same window settings you used in part A, explore the graphs of the following rational functions. Sketch each graph on separate axes, and note any holes or asymptotes.
- i) $f(x) = \frac{x^2 - 1}{x - 1}$
 - ii) $f(x) = \frac{3}{x + 1}$
 - iii) $f(x) = \frac{x + 1}{x^2 - 2x - 3}$
 - iv) $f(x) = \frac{x + 1}{x + 2}$
 - v) $f(x) = \frac{0.5x^2 + 1}{x - 1}$
 - vi) $f(x) = \frac{x^2 + 2x}{x + 1}$
 - vii) $f(x) = \frac{9x}{1 + x^2}$
 - viii) $f(x) = \frac{2x^2 - 3}{x^2 + 1}$
- G.** Examine the graphs of the functions in parts i) and v) of part F at the point where $x = 1$. Explain why $f(x) = \frac{x^2 - 1}{x - 1}$ has a hole where $x = 1$, but $f(x) = \frac{0.5x^2 + 1}{x - 1}$ has a vertical asymptote. Identify the other functions in part F that have holes and the other functions that have vertical asymptotes.

Tech Support

When entering a rational function into a graphing calculator, use brackets around the expression in the numerator and the expression in the denominator.

oblique asymptote

an asymptote that is neither vertical nor horizontal, but slanted



- H. Redraw the graph of the rational function $f(x) = \frac{0.5x^2 + 1}{x - 1}$. Then enter the equation $y = 0.5x + 0.5$ into Y2 of the equation editor. What do you notice? Examine all your other sketches in this exploration to see if any of the other functions have an oblique asymptote.
- I. Examine the equations with graphs that have horizontal asymptotes in part F. Compare the degree of the expression in the numerator with the degree of the expression in the denominator. Is there a connection between the degrees in the numerator and denominator and the existence of horizontal asymptotes? Explain. Repeat for functions with oblique asymptotes.
- J. Investigate several functions of the form $f(x) = \frac{ax + b}{cx + d}$. Note similarities and differences. Without graphing, how can you predict where a horizontal asymptote will occur?
- K. Investigate graphs of quotients of quadratic functions. How are they different from graphs of quotients of linear functions?
- L. Summarize the different characteristics of the graphs of rational functions.

Reflecting

- M. How do the zeros of the function in the numerator help you graph the rational function? How do the zeros of the function in the denominator help you graph the rational function?
- N. Explain how you can use the expressions in the numerator and the denominator of a rational function to decide if the graph has
- a hole
 - a vertical asymptote
 - a horizontal asymptote
 - an oblique asymptote

In Summary


Key Ideas

- The quotient of two polynomial functions results in a rational function which often has one or more discontinuities.
- The breaks or discontinuities in a rational function occur where the function is undefined. The function is undefined at values where the denominator is equal to zero. As a result, these values must be restricted from the domain of the function.
- The values that must be restricted from the domain of a rational function result in key characteristics that define the shape of the graph. These characteristics include a combination of vertical asymptotes (also called infinite discontinuities) and holes (also called point discontinuities).
- The end behaviours of many rational functions are determined by either horizontal asymptotes or oblique asymptotes.

Need to Know


- A rational function, $f(x) = \frac{p(x)}{q(x)}$, has a hole at $x = a$ if $\frac{p(a)}{q(a)} = \frac{0}{0}$. This occurs when $p(x)$ and $q(x)$ contain a common factor of $(x - a)$.

For example, $f(x) = \frac{x^2 - 4}{x - 2}$ has the common factor of $(x - 2)$ in the numerator and the denominator. This results in a hole in the graph of $f(x)$ at $x = 2$.




- A rational function, $f(x) = \frac{p(x)}{q(x)}$, has a vertical asymptote at $x = a$ if $\frac{p(a)}{q(a)} = \frac{p(a)}{0}$.

For example, $f(x) = \frac{x+1}{x-2}$ has a vertical asymptote at $x = 2$.




- A rational function, $f(x) = \frac{p(x)}{q(x)}$, has a horizontal asymptote only when the degree of $p(x)$ is less than or equal to the degree of $q(x)$. For example,

$f(x) = \frac{2x}{x+1}$ has a horizontal asymptote at $y = 2$.



- A rational function, $f(x) = \frac{p(x)}{q(x)}$, has an oblique (slant) asymptote only when the degree of $p(x)$ is greater than the degree of $q(x)$ by exactly 1. For example,

$f(x) = \frac{x^2 + 4}{x + 1}$ has an oblique asymptote.



FURTHER Your Understanding

1. Without using graphing technology, match each equation with its corresponding graph. Explain your reasoning.

a) $y = \frac{-1}{x-3}$

d) $y = \frac{x}{(x-1)(x+3)}$


b) $y = \frac{x^2-9}{x-3}$

e) $y = \frac{1}{x^2+5}$


c) $y = \frac{1}{(x+3)^2}$

f) $y = \frac{x^2}{x-3}$


A




C




E




B



D



F



2. For each function, determine the equations of any vertical asymptotes, the locations of any holes, and the existence of any horizontal or oblique asymptotes.

a) $y = \frac{x}{x+4}$

e) $y = \frac{1}{(x+3)(x-5)}$

i) $y = \frac{8x}{4x+1}$

b) $y = \frac{1}{2x+3}$

f) $y = \frac{-x}{x+1}$

j) $y = \frac{x+4}{x^2-16}$

c) $y = \frac{2x+5}{x-6}$

g) $y = \frac{3x-6}{x-2}$

k) $y = \frac{x}{5x-3}$

d) $y = \frac{x^2-9}{x+3}$

h) $y = \frac{-4x+1}{2x-5}$

l) $y = \frac{-3x+1}{2x-8}$

3. Write an equation for a rational function with the properties as given.

- a hole at $x = 1$
- a vertical asymptote anywhere and a horizontal asymptote along the x -axis
- a hole at $x = -2$ and a vertical asymptote at $x = 1$
- a vertical asymptote at $x = -1$ and a horizontal asymptote at $y = 2$
- an oblique asymptote, but no vertical asymptote

5.3

Graphs of Rational Functions of the Form $f(x) = \frac{ax + b}{cx + d}$

GOAL

Sketch the graphs of rational functions, given equations of the form $f(x) = \frac{ax + b}{cx + d}$.

YOU WILL NEED

- graph paper
- graphing calculator or graphing software

INVESTIGATE the Math



The radius, in centimetres, of a circular juice blot on a piece of paper towel is modelled by $r(t) = \frac{1 + 2t}{1 + t}$, where t is measured in seconds. According to this model, the maximum size of the blot is determined by the location of the horizontal asymptote.

? How can you find the equation of the horizontal asymptote of a rational function of the form $f(x) = \frac{ax + b}{cx + d}$?

- A.** Without graphing, determine the domain, intercepts, vertical asymptote, and positive/negative intervals of the simple rational function $f(x) = \frac{x}{x + 1}$.
- B.** Copy the following tables, and complete them by evaluating $f(x)$ for each value of x . Examine the **end behaviour** of $f(x)$ by observing the trend in $f(x)$ as x grows positively large and negatively large. What value does $f(x)$ seem to approach?

$x \rightarrow \infty$	
x	$f(x) = \frac{x}{x + 1}$
10	
100	
1 000	
10 000	
100 000	
1 000 000	

$x \rightarrow -\infty$	
x	$f(x) = \frac{x}{x + 1}$
-10	
-100	
-1 000	
-10 000	
-100 000	
-1 000 000	

- C. Write an equation for the horizontal asymptote of the function in part B.
- D. Repeat parts A, B, and C for the functions $g(x) = \frac{4x}{x+1}$, $h(x) = \frac{2x}{3x+1}$, and $m(x) = \frac{3x-2}{2x-5}$.
- E. Verify your results by graphing all the functions in part D on a graphing calculator. Note similarities and differences among the graphs.
- F. Make a list of the equations of the functions and the equations of their horizontal asymptotes. Discuss how the degree of the numerator compares with the degree of the denominator. Explain how the leading coefficients of x in the numerator and the denominator determine the equation of the horizontal asymptote.
- G. Determine the equation of the horizontal asymptote of the juice blot function $r(t) = \frac{1+2t}{1+t}$. What does this equation tell you about the eventual size of the juice blot?

Reflecting

- H. How do the graphs of rational functions with linear expressions in the numerator and denominator compare with the graphs of reciprocal functions?
- I. Explain how you determined the equation of a horizontal asymptote from
- end behaviour tables
 - the equation of the function

APPLY the Math

EXAMPLE 1

Selecting a strategy to determine how a graph approaches a vertical asymptote

Determine how the graph of $f(x) = \frac{3x-5}{x+2}$ approaches its vertical asymptote.

Solution

$f(x) = \frac{3x-5}{x+2}$ has a vertical asymptote with the equation $x = -2$. Near this asymptote, the values of the function will grow very large in a positive direction or very large in a negative direction.

$f(x)$ is undefined when $x = -2$.
There is no common factor in the numerator and denominator.

Choose a value of x to the left and very close to -2 . This value is less than -2 .

$$\begin{aligned} f(-2.1) &= \frac{3(-2.1) - 5}{(-2.1) + 2} \\ &= 113 \end{aligned}$$

The graph of a rational function never crosses a vertical asymptote, so choose x -values that are very close to the vertical asymptote, on both sides, to determine the behaviour of the function.

On the left side of the vertical asymptote, the values of the function are positive. As $x \rightarrow -2$, $f(x) \rightarrow \infty$.


The function increases to large positive values as x approaches -2 from the left.

Choose a value of x to the right and very close to -2 . This value is greater than -2 .

$$\begin{aligned} f(-1.9) &= \frac{3(-1.9) - 5}{(-1.9) + 2} \\ &= -107 \end{aligned}$$

On the right side of the vertical asymptote, the values of the function are negative. As $x \rightarrow -2$, $f(x) \rightarrow -\infty$.

The function decreases to small negative values as x approaches -2 from the right.



Make a sketch to show how the graph approaches the vertical asymptote.

EXAMPLE 2

Using key characteristics to sketch the graph of a rational function

For each function,

a) $f(x) = \frac{2}{x-3}$

b) $f(x) = \frac{x-2}{3x+4}$

c) $f(x) = \frac{x-3}{2x-6}$

- i) determine the domain, intercepts, asymptotes, and positive/negative intervals
- ii) use these characteristics to sketch the graph of the function
- iii) describe where the function is increasing or decreasing



Solution

a) $f(x) = \frac{2}{x-3}$

i) $D = \{x \in \mathbf{R} | x \neq 3\}$

The function $f(x) = \frac{2}{x-3}$ is undefined when $x = 3$.

$f(0) = -\frac{2}{3}$, so the y -intercept is $-\frac{2}{3}$.

$f(x) \neq 0$, so there is no x -intercept.

Any rational function equals zero when its numerator equals zero. The numerator is always 2, so $f(x)$ can never equal zero.

The line $x = 3$ is a vertical asymptote.
The line $y = 0$ is a horizontal asymptote.

Since the numerator and denominator do not contain the common factor $(x - 3)$, $f(x)$ has a vertical asymptote at $x = 3$. Any rational function that is formed by a constant numerator and a linear function denominator has a horizontal asymptote at $y = 0$.

$f(x)$ is negative when $x \in (-\infty, 3)$ and positive when $x \in (3, \infty)$.

The numerator is always positive, so the denominator determines the sign of $f(x)$.
 $x - 3 < 0$ when $x < 3$
 $x - 3 > 0$ when $x > 3$

ii) Confirm the behaviour of $f(x)$ near the vertical asymptote.


$f(3.1) = 20$, so as

$x \rightarrow 3, f(x) \rightarrow \infty$

on the right.

$f(2.9) = -20$, so as

$x \rightarrow 3, f(x) \rightarrow -\infty$ on the left.



Use all the information in part i) to sketch the graph.

iii) From the graph, the function is decreasing on its entire domain: when $x \in (-\infty, 3)$ and when $x \in (3, \infty)$.

b) $f(x) = \frac{x-2}{3x+4}$

i) $3x + 4 \neq 0$
 $3x \neq -4$

$$x \neq -\frac{4}{3}$$

$$D = \left\{x \in \mathbf{R} \mid x \neq -\frac{4}{3}\right\}$$

$f(x)$ is undefined when the denominator is zero.

$$f(0) = \frac{0-2}{3(0)+4} = \frac{-2}{4} \text{ or } -\frac{1}{2}$$

To determine the y -intercept, let $x = 0$.

The y -intercept is $-\frac{1}{2}$.

$$f(x) = 0 \text{ when } \frac{x-2}{3x+4} = 0.$$

$$x-2 = 0$$

$$x = 2$$

To determine the x -intercept, let $y = 0$. Any rational function equals zero when its numerator equals zero.

The x -intercept is 2.

The line $x = -\frac{4}{3}$ is a vertical asymptote.

This is the value that makes $f(x)$ undefined.

The line $y = \frac{1}{3}$ is a horizontal asymptote.

The ratio of the leading coefficients of the numerator and denominator is $\frac{1}{3}$.

Examine the signs of the numerator and denominator, and their quotient, to determine the positive/negative intervals.


	$x < -\frac{4}{3}$	$-\frac{4}{3} < x < 2$	$x > 2$
$x - 2$	-	-	+
$3x + 4$	-	+	+
$\frac{x-2}{3x+4}$	$\frac{-}{-} = +$	$\frac{-}{+} = -$	$\frac{+}{+} = +$

The vertical asymptote and the x -intercept divide the set of real numbers into three intervals: $(-\infty, -\frac{4}{3})$, $(-\frac{4}{3}, 2)$, and $(2, \infty)$. Choose numbers in each interval to evaluate the sign of each expression.

$f(x)$ is positive when $x \in (-\infty, -\frac{4}{3})$
 and when $x \in (2, \infty)$.

$f(x)$ is negative when $x \in (-\frac{4}{3}, 2)$.





When sketching the graph, it helps to shade the regions where there is no graph. Use the positive and negative intervals as indicators for these regions. For example, since $f(x)$ is positive on $(-\infty, -\frac{4}{3})$, there is no graph under the x -axis on this interval. Draw the asymptotes, and mark the intercepts. Then draw the graph to approach the asymptotes.

- iii) From the graph, $f(x)$ is increasing on its entire domain; that is, when $x \in (-\infty, -\frac{4}{3})$ and when $x \in (-\frac{4}{3}, \infty)$.

c) $f(x) = \frac{x-3}{2x-6}$

i) $2x - 6 \neq 0$
 $2x \neq 6$
 $x \neq 3$

$f(x)$ is undefined when the denominator is zero.

$D = \{x \in \mathbf{R} | x \neq 3\}$

$f(0) = \frac{0-3}{2(0)-6} = \frac{-3}{-6}$ or $\frac{1}{2}$

To determine the y -intercept, let $x = 0$.

The y -intercept is $\frac{1}{2}$.

$f(x) \neq 0$, so there is no x -intercept.

To determine the x -intercept, let $y = 0$. Only consider when the numerator is zero; that is, when $x - 3 = 0$. Therefore, the numerator is zero at $x = 3$, but this has already been excluded from the domain.

$f(x) = \frac{x-3}{2(x-3)}$

$f(x)$ has a hole, not a vertical asymptote, where $x = 3$.

Factoring reveals a common factor $(x - 3)$ in the numerator and denominator. The graph has a hole at the point where $x = 3$.


$f(x) = \frac{\overset{1}{x-3}}{2(\overset{1}{x-3})} = \frac{1}{2}$

The value of the function is always $\frac{1}{2}$ for all values of x , except when $x = 3$.

$f(x)$ is positive at all points in its domain.

$f(x)$ has no vertical asymptote or x -intercept. There is only one interval to consider: $(-\infty, \infty)$.
For any value of x , $f(x) = \frac{1}{2}$.

- ii) The graph is a horizontal line with the equation $y = \frac{1}{2}$. There is a hole at $x = 3$.



Use the information in part i) to sketch the graph.

- iii) The function is neither increasing nor decreasing. It is constant on its entire domain.

EXAMPLE 3

Solving a problem by graphing a rational function

The function $P(t) = \frac{30(7t + 9)}{3t + 2}$ models the population, in thousands, of a town t years since 1990. Describe the population of the town over the next 20 years.


Solution

$$P(t) = \frac{30(7t + 9)}{3t + 2}$$


Determine the initial population in 1990, when $t = 0$.

$$P(0) = \frac{30(7(0) + 9)}{3(0) + 2} = 135$$


Use the equation to help you decide on the window settings. For the given context, $t \geq 0$ and $P(t) > 0$.



Graph $P(t)$ to show the population for the 20 years after 1990.




The value that makes $3t + 2 = 0$ lies outside the domain of $P(t)$. There is no vertical asymptote in the domain.



TRACE along the curve to get an idea of how the population changed.

In the first two years, the population dropped by about 50 000 people. Then it began to level off and approach a steady value.

There is a horizontal asymptote at $P = \frac{30(7)}{3} = 70$.



Use the function equation to determine the equation of the horizontal asymptote. For large values of t , $P(t) \doteq \frac{30(7t)}{3t}$. Therefore, the leading coefficients in the numerator and denominator define the equation of the horizontal asymptote.

The population of the town has been decreasing since 1990. It was 135 000 in 1990, but dropped by about 50 000 in the next two years. Since then, the population has begun to level off and, according to the model, will approach a steady value of 70 000 people by 2010.

Multiply the values of the function by 1000, since the population is given in thousands.


In Summary

Key Ideas


- The graphs of most rational functions of the form $f(x) = \frac{b}{cx + d}$ and $f(x) = \frac{ax + b}{cx + d}$ have both a vertical asymptote and a horizontal asymptote.
- You can determine the equation of the vertical asymptote directly from the equation of the function by finding the zero of the denominator.
- You can determine the equation of the horizontal asymptote directly from the equation of the function by examining the ratio of the leading coefficients in the numerator and the denominator. This gives you the end behaviours of the function.
- To sketch the graph of a rational function, you can use the domain, intercepts, equations of asymptotes, and positive/negative intervals.

Need to Know

- Rational functions of the form $f(x) = \frac{b}{cx + d}$ have a vertical asymptote defined by $x = -\frac{d}{c}$ and a horizontal asymptote defined by $y = 0$. For example, see the graph of $f(x) = \frac{2}{x-3}$.




- Most rational functions of the form $f(x) = \frac{ax + b}{cx + d}$ have a vertical asymptote defined by $x = -\frac{d}{c}$ and a horizontal asymptote defined by $y = \frac{a}{c}$. For example, see the graph of $f(x) = \frac{4x-1}{2x-1}$.



The exception occurs when the numerator and the denominator both contain a common linear factor. This results in a graph of a horizontal line that has a hole where the zero of the common factor occurs. As a result, the graph has no asymptotes. For example, see the

graph of $f(x) = \frac{4x-8}{x-2} = \frac{4(x-2)}{(x-2)}$.




CHECK Your Understanding


1. Match each function with its graph.

a) $h(x) = \frac{x+4}{2x+5}$ c) $f(x) = \frac{3}{x-1}$
 b) $m(x) = \frac{2x-4}{x-2}$ d) $g(x) = \frac{2x-3}{x+2}$


A




C



B



D



2. Consider the function $f(x) = \frac{3}{x-2}$.

- State the equation of the vertical asymptote.
- Use a table of values to determine the behaviour(s) of the function near its vertical asymptote.
- State the equation of the horizontal asymptote.
- Use a table of values to determine the end behaviours of the function near its horizontal asymptote.
- Determine the domain and range.
- Determine the positive and negative intervals.
- Sketch the graph.

3. Repeat question 2 for the rational function $f(x) = \frac{4x-3}{x+1}$.

PRACTISING

4. State the equation of the vertical asymptote of each function. Then choose a strategy to determine how the graph of the function approaches its vertical asymptote.
- a) $y = \frac{2}{x+3}$ c) $y = \frac{2x+1}{2x-1}$
 b) $y = \frac{x-1}{x-5}$ d) $y = \frac{3x+9}{4x+1}$
5. For each function, determine the domain, intercepts, asymptotes, and **K** positive/negative intervals. Use these characteristics to sketch the graph of the function. Then describe where the function is increasing or decreasing.
- a) $f(x) = \frac{3}{x+5}$ c) $f(x) = \frac{x+5}{4x-1}$
 b) $f(x) = \frac{10}{2x-5}$ d) $f(x) = \frac{x+2}{5(x+2)}$
6. Read each set of conditions. State the equation of a rational function of the form $f(x) = \frac{ax+b}{cx+d}$ that meets these conditions, and sketch the graph.
- a) vertical asymptote at $x = -2$, horizontal asymptote at $y = 0$; negative when $x \in (-\infty, -2)$, positive when $x \in (-2, \infty)$; always decreasing
 b) vertical asymptote at $x = -2$, horizontal asymptote at $y = 1$; x -intercept = 0, y -intercept = 0; positive when $x \in (-\infty, -2)$ or $(0, \infty)$, negative when $x \in (-2, 0)$
 c) hole at $x = 3$; no vertical asymptotes; y -intercept = $(0, 0.5)$
 d) vertical asymptotes at $x = -2$ and $x = 6$, horizontal asymptote at $y = 0$; positive when $x \in (-\infty, -2)$ or $(6, \infty)$, negative when $x \in (-2, 6)$; increasing when $x \in (-\infty, 2)$, decreasing when $x \in (2, \infty)$
7. a) Use a graphing calculator to investigate the similarities and **T** differences in the graphs of rational functions of the form $f(x) = \frac{8x}{nx+1}$, for $n = 1, 2, 4$, and 8.
 b) Use your answer for part a) to make a conjecture about how the function changes as the values of n approach infinity.
 c) If n is negative, how does the function change as the value of n approaches negative infinity? Choose your own values, and use them as examples to support your conclusions.

8. Without using a graphing calculator, compare the graphs of the rational functions $f(x) = \frac{3x+4}{x-1}$ and $g(x) = \frac{x-1}{2x+3}$.
9. The function $I(t) = \frac{15t+25}{t}$ gives the value of an investment, in thousands of dollars, over t years.
- What is the value of the investment after 2 years?
 - What is the value of the investment after 1 year?
 - What is the value of the investment after 6 months?
 - There is an asymptote on the graph of the function at $t = 0$. Does this make sense? Explain why or why not.
 - Choose a very small value of t (a value near zero). Calculate the value of the investment at this time. Do you think that the function is accurate at this time? Why or why not?
 - As time passes, what will the value of the investment approach?
10. An amount of chlorine is added to a swimming pool that contains pure water. The concentration of chlorine, c , in the pool at t hours is given by $c(t) = \frac{2t}{2+t}$, where c is measured in milligrams per litre. What happens to the concentration of chlorine in the pool during the 24 h period after the chlorine is added?
11. Describe the key characteristics of the graphs of rational functions of the form $f(x) = \frac{ax+b}{cx+d}$. Explain how you can determine these characteristics using the equations of the functions. In what ways are the graphs of all the functions in this family alike? In what ways are they different? Use examples in your comparison.

Extending

12. Not all asymptotes are horizontal or vertical. Find a rational function that has an asymptote that is neither horizontal nor vertical, but slanted or oblique.
13. Use long division to rewrite $f(x) = \frac{2x^3 - 7x^2 + 8x - 5}{x - 1}$ in the form $f(x) = ax^2 + bx + c + \frac{k}{x-1}$. What does this tell you about the end behaviour of the function? Graph the function. Include all asymptotes in your graph. Write the equations of the asymptotes.
14. Let $f(x) = \frac{3x-1}{x^2-2x-3}$, $g(x) = \frac{x^3+8}{x^2+9}$, $h(x) = \frac{x^3-3x}{x+1}$, and $m(x) = \frac{x^2+x-12}{x^2-4}$.
- Which of these rational functions has a horizontal asymptote?
 - Which has an oblique asymptote?
 - Which has no vertical asymptote?
 - Graph $y = m(x)$, showing the asymptotes and intercepts.

5.4

Solving Rational Equations

YOU WILL NEED

- graphing calculator or graphing software

GOAL

Connect the solution to a rational equation with the graph of a rational function.

LEARN ABOUT the Math



When they work together, Stuart and Lucy can deliver flyers to all the homes in their neighbourhood in 42 min. When Lucy works alone, she can finish the deliveries in 13 min less time than Stuart can when he works alone.

- ? When Stuart works alone, how long does he take to deliver the flyers?

EXAMPLE 1 Selecting a strategy to solve a rational equation

Determine the time that Stuart takes to deliver the flyers when he works alone.

Solution A: Creating an equation and solving it using algebra

Let s minutes be the time that Stuart takes to deliver the flyers when working alone.

Lucy takes $(s - 13)$ minutes when working alone.

Choose a variable to represent Stuart's time and use it to write an expression for Lucy's time.

Lucy delivers the flyers in 13 min less time than Stuart.

The fraction of deliveries made in one minute

- by Stuart working alone is $\frac{1}{s}$
- by Lucy working alone is $\frac{1}{s - 13}$
- by Stuart and Lucy working together is $\frac{1}{42}$

Compare the rates at which they work.

For example, if Stuart took 80 min to deliver all the flyers, he would deliver $\frac{1}{80}$ of the flyers per minute.

$s > 13$ because Stuart takes longer than Lucy to deliver the flyers, and it is not possible for the denominators to be zero.

$$\frac{1}{s} + \frac{1}{s - 13} = \frac{1}{42}$$



Multiply by the LCD.

$$42s(s-13)\left(\frac{1}{s} + \frac{1}{s-13}\right) = 42s(s-13)\left(\frac{1}{42}\right)$$

$$\frac{42s(s-13)}{s} + \frac{42s(s-13)}{s-13} = \frac{42s(s-13)}{42}$$

$$\frac{42\cancel{s}(s-13)}{\cancel{s}} + \frac{42\cancel{s}(\cancel{s-13})}{\cancel{s-13}} = \frac{42\cancel{s}(s-13)}{\cancel{42}}$$

$$42(s-13) + 42s = s(s-13)$$

There are no common factors in the denominators, so the LCD (lowest common denominator) is the product of the three denominators $42s(s-13)$. Multiply each term by the LCD, and then simplify the resulting rational expressions to remove all the denominators.

$$42s - 546 + 42s = s^2 - 13s$$

$$0 = s^2 - 97s + 546$$

$$0 = (s-91)(s-6)$$

$$s = 6 \text{ or } 91$$

Solve the quadratic equation by factoring or by using the quadratic formula.

$s > 13$ so 6 is not an admissible solution.

Remember to look for inadmissible solutions by carefully considering both the context and the information given in the problem.

$$\text{LS} = \frac{1}{s} + \frac{1}{s-13}$$

$$= \frac{1}{91} + \frac{1}{91-13}$$

$$= \frac{1}{91} + \frac{1}{78}$$

$$= \frac{78+91}{7098}$$

$$= \frac{169}{7098}$$

$$= \frac{1}{42}$$

$$\text{RS} = \frac{1}{42}$$

Check the solution, $s = 91$, by substituting it into the original equation.

Since $\text{LS} = \text{RS}$, $s = 91$ is the solution.

It will take Stuart 91 min to deliver the flyers when working alone.



Solution B: Using the graph of a rational function to solve a rational equation

The equation that models the problem is $\frac{1}{s} + \frac{1}{s-13} = \frac{1}{42}$, where s represents the time, in minutes, that Stuart takes to deliver the flyers when working alone.

$$\frac{1}{s} + \frac{1}{s-13} - \frac{1}{42} = 0$$

Subtract $\frac{1}{42}$ from each side.

To solve the equation, find the zeros of the function

$$f(s) = \frac{1}{s} + \frac{1}{s-13} - \frac{1}{42}.$$



$$\text{Graph } f(s) = \frac{1}{s} + \frac{1}{s-13} - \frac{1}{42}.$$

From the equation, you can expect the graph to have vertical asymptotes at $s = 0$ and $s = 13$.

Since you are only interested in finding the zeros, you can limit the y -values to those close to zero.

Tech Support

For help determining the zeros on a graphing calculator, see Technical Appendix, T-8.




The first zero for $f(s)$ is $s = 6$.
Reject this solution since $s > 13$.

Determine the other zero.

You know that Lucy takes 13 min less time than Stuart takes, so Stuart must take longer than 13 min.

The solution $s = 6$ is inadmissible.



The solution is $s = 91$.

Stuart takes 91 min to deliver the flyers when working alone.

Reflecting

- A. In Solution A, explain how a rational equation was created using the times given in the problem.
- B. In Solution B, explain how finding the zeros of a rational function provided the solution to the problem.
- C. Where did the inadmissible root obtained in Solution A show up in the graphical solution in Solution B? How was this root dealt with?

APPLY the Math

EXAMPLE 2

Using an algebraic strategy to solve simple rational equations

Solve each rational equation.

a) $\frac{x-2}{x-3} = 0$ b) $\frac{x+3}{x-4} = \frac{x-1}{x+2}$

Solution

a) $\frac{x-2}{x-3} = 0, x \neq 3$ ← Determine any restrictions on the value of x .

$$\begin{aligned} \cancel{(x-3)}^1 \frac{(x-2)}{\cancel{(x-3)}^1} &= 0(x-3) \\ x-2 &= 0 \\ x &= 2 \end{aligned}$$

← Multiply both sides of the equation by the LCD, $(x-3)$.
← Add 2 to each side.

To verify, graph $f(x) = \frac{x-2}{x-3}$

and use the zero operation to determine the zero.

← From the equation, the graph will have a vertical asymptote at $x = 3$ and a horizontal asymptote at $y = 1$.



The solution is $x = 2$.

b) $\frac{x+3}{x-4} = \frac{x-1}{x+2}, x \neq -2, 4$ ← Note the restrictions.

$$(x-4)(x+2)\left(\frac{x+3}{x-4}\right) = (x-4)(x+2)\left(\frac{x-1}{x+2}\right)$$

← Multiply each side of the equation by the LCD, $(x-4)(x+2)$.

$$\cancel{(x-4)}^1(x+2)\left(\frac{x+3}{\cancel{x-4}^1}\right) = (x-4)\cancel{(x+2)}^1\left(\frac{x-1}{\cancel{x+2}^1}\right)$$

← Simplify.

$$(x+2)(x+3) = (x-4)(x-1)$$

$$x^2 + 5x + 6 = x^2 - 5x + 4$$

← Expand. Subtract x^2 , and add $5x$ to both sides.



$$10x + 6 = 4$$

← Solve the resulting linear equation.

$$10x = -2$$

$$x = -0.2$$

To verify, graph $f(x) = \frac{x+3}{x-4} - \frac{x-1}{x+2}$ and use the zero operation to determine the zero.

← Adjust the window settings so you can view enough of the graph to see all the possible zeros.

The solution is $x = -0.2$.

EXAMPLE 3

Connecting the solution to a problem with the zeros of a rational function

Salt water is flowing into a large tank that contains pure water. The concentration of salt, c , in the tank at t minutes is given by $c(t) = \frac{10t}{25+t}$, where c is measured in grams per litre. When does the salt concentration in the tank reach 3.75 g/L?

Solution

If the salt concentration is 3.75, $c(t) = 3.75$.

$$\frac{10t}{25+t} = 3.75$$

$$(25+t)\left(\frac{10t}{25+t}\right) = 3.75(25+t)$$

$$\cancel{(25+t)}^1 \frac{10t}{\cancel{(25+t)}^1} = 3.75(25+t)$$

$$10t = 93.75 + 3.75t$$

← Set the function expression equal to 3.75. $25+t \neq 0$, and because t measures the time since the salt water started flowing, $t \geq 0$.

← Multiply both sides of the equation by the LCD, $(25+t)$, and solve the resulting linear equation.

$$10t - 3.75t = 93.75$$

$$6.25t = 93.75$$

$$\frac{6.25t}{6.25} = \frac{93.75}{6.25}$$


$$t = 15$$

Use inverse operations to solve for t .


It takes 15 min for the salt concentration to reach 3.75 g/L.

To verify, graph $f(t) = \frac{10t}{25 + t}$ and $g(t) = 3.75$, and determine where the functions intersect.

Use an appropriate window setting, based on the domain, $t \geq 0$.



The salt concentration reaches 3.75 g/L after 15 min.



Use the intersect operation.

Tech Support

For help determining the point of intersection between two functions, see Technical Appendix, T-12.

EXAMPLE 4 Using a rational function to model and solve a problem

Rima bought a case of concert T-shirts for \$450. She kept two T-shirts for herself and sold the rest for \$560, making a profit of \$10 on each T-shirt. How many T-shirts were in the case?

Solution

Let the number of T-shirts in the case be x .

$$\text{Buying price per T-shirt} = \frac{450}{x}$$

$$\text{Selling price per T-shirt} = \frac{560}{x - 2}$$

Rima paid \$450 for x T-shirts, so each T-shirt cost her $\$ \frac{450}{x}$.

She kept two for herself, which left $x - 2$ T-shirts for her to sell.

Rima sold $x - 2$ T-shirts for \$560, so she charged $\$ \frac{560}{x - 2}$ for each one.

$$\frac{560}{x-2} - \frac{450}{x} = 10$$

She made a profit of \$10 on each T-shirt, so the difference between the selling price and the buying price was \$10.

$$x(x-2)\left(\frac{560}{x-2} - \frac{450}{x}\right) = 10x(x-2)$$

Multiply both sides of the equation by the LCD, $x(x-2)$.

$$\frac{560x\cancel{(x-2)}}{\cancel{x-2}} - \frac{450\cancel{x}(x-2)}{\cancel{x}} = 10x(x-2)$$

$$560x - 450(x-2) = 10x(x-2)$$

Expand and collect all terms to one side of the equation.

$$560x - 450x + 900 = 10x^2 - 20x$$

$$0 = 10x^2 - 130x - 900$$

$$0 = 10(x^2 - 13x - 90)$$

Solve the resulting quadratic equation by factoring.

$$0 = 10(x-18)(x+5)$$


$$x = 18 \text{ or } -5$$

-5 is inadmissible since $x \geq 0$.
There were 18 T-shirts in the case.


You cannot have a negative number of T-shirts in the case.

To verify, graph $f(x) = \frac{560}{x-2} - \frac{450}{x} - 10$ and determine the zeros using the zero operation.


If $\frac{560}{x-2} - \frac{450}{x} = 10$, then $\frac{560}{x-2} - \frac{450}{x} - 10 = 0$.
Zeros for $f(x)$ are possible solutions to the problem.



Use an appropriate window setting, based on the domain, $x \geq 0$.



The zero occurs when $x = 18$.
Zoom out to check that there are no other zeros in the domain.



The other zero is for a negative value of x , which is inadmissible in the context of this problem.

There is no other zero in the domain.
There were 18 T-shirts in the case.

In Summary

Key Ideas

- You can solve a rational equation algebraically by multiplying each term in the equation by the lowest common denominator and solving the resulting polynomial equation.
- The root of the equation $\frac{ax + b}{cx + d} = 0$ is the zero (x-intercept) of the function $f(x) = \frac{ax + b}{cx + d}$.
- You can use graphing technology to solve a rational equation or verify the solution. Determine the zeros of the corresponding rational function, or determine the intersection of two functions.

Need to Know

- The zeros of a rational function are the zeros of the function in the numerator.
- Reciprocal functions do not have zeros. All functions of the form $f(x) = \frac{1}{g(x)}$ have the x-axis as a horizontal asymptote. They do not intersect the x-axis.
- When solving contextual problems, it is important to check for inadmissible solutions that are outside the domain determined by the context.
- When using a graphing calculator to determine a zero or intersection point, you can avoid inadmissible roots by matching the window settings to the domain of the function in the context of the problem.

CHECK Your Understanding

- Are $x = 3$ and $x = -2$ solutions to the equation $\frac{2}{x} = \frac{x-1}{3}$? Explain how you know.
- Solve each equation algebraically. Then verify your solution using graphing technology.

a) $\frac{x+3}{x-1} = 0$	c) $\frac{x+3}{x-1} = 2x+1$
b) $\frac{x+3}{x-1} = 2$	d) $\frac{3}{3x+2} = \frac{6}{5x}$
- For each rational equation, write a function whose zeros are the solutions.

a) $\frac{x-3}{x+3} = 2$	c) $\frac{x-1}{x} = \frac{x+1}{x+3}$
b) $\frac{3x-1}{x} = \frac{5}{2}$	d) $\frac{x-2}{x+3} = \frac{x-4}{x+5}$
- Solve each equation in question 3 algebraically, and verify your solution using a graphing calculator.

PRACTISING

5. Solve each equation algebraically.

a) $\frac{2}{x} + \frac{5}{3} = \frac{7}{x}$

d) $\frac{2}{x+1} + \frac{1}{x+1} = 3$

b) $\frac{10}{x+3} + \frac{10}{3} = 6$

e) $\frac{2}{2x+1} = \frac{5}{4-x}$

c) $\frac{2x}{x-3} = 1 - \frac{6}{x-3}$

f) $\frac{5}{x-2} = \frac{4}{x+3}$

6. Solve each equation algebraically.

a) $\frac{2x}{2x+1} = \frac{5}{4-x}$

d) $x + \frac{x}{x-2} = 0$

b) $\frac{3}{x} + \frac{4}{x+1} = 2$

e) $\frac{1}{x+2} + \frac{24}{x+3} = 13$

c) $\frac{2x}{5} = \frac{x^2 - 5x}{5x}$

f) $\frac{-2}{x-1} = \frac{x-8}{x+1}$

7. Solve each equation using graphing technology. Round your answers to two decimal places, if necessary.

a) $\frac{2}{x+2} = \frac{3}{x+6}$

d) $\frac{1}{x} - \frac{1}{45} = \frac{1}{2x-3}$

b) $\frac{2x-5}{x+10} = \frac{1}{x-6}$

e) $\frac{2x+3}{3x-1} = \frac{x+2}{4}$

c) $\frac{1}{x-3} = \frac{x+2}{7x+14}$

f) $\frac{1}{x} = \frac{2}{x} + 1 + \frac{1}{1-x}$

8. a) Use algebra to solve $\frac{x+1}{x-2} = \frac{x+3}{x-4}$. Explain your steps.

K b) Verify your answer in part a) using substitution.

c) Verify your answer in part a) using a graphing calculator.

9. The Greek mathematician Pythagoras is credited with the discovery of the Golden Rectangle. This is considered to be the rectangle with the dimensions that are the most visually appealing. In a Golden Rectangle, the length and width are related by the proportion

$$\frac{l}{w} = \frac{w}{l-w}$$

A billboard with a length of 15 m is going to be built.

What must its width be to form a Golden Rectangle?

10. The Turtledove Chocolate factory has two chocolate machines. Machine A takes s minutes to fill a case with chocolates, and machine B takes $s + 10$ minutes to fill a case. Working together, the two machines take 15 min to fill a case. Approximately how long does each machine take to fill a case?

11. Tayla purchased a large box of comic books for \$300. She gave 15 of the comic books to her brother and then sold the rest on an Internet website for \$330, making a profit of \$1.50 on each one. How many comic books were in the box? What was the original price of each comic book?
12. Polluted water flows into a pond. The concentration of pollutant, c , in the pond at time t minutes is modelled by the equation $c(t) = 9 - 90\,000\left(\frac{1}{10\,000 + 3t}\right)$, where c is measured in kilograms per cubic metre.
- When will the concentration of pollutant in the pond reach 6 kg/m^3 ?
 - What will happen to the concentration of pollutant over time?
13. Three employees work at a shipping warehouse. Tom can fill an order in s minutes. Paco can fill an order in $s - 2$ minutes. Carl can fill an order in $s + 1$ minutes. When Tom and Paco work together, they take about 1 minute and 20 seconds to fill an order. When Paco and Carl work together, they take about 1 minute and 30 seconds to fill an order.
- How long does each person take to fill an order?
 - How long would all three of them, working together, take to fill an order?
14. Compare and contrast the different methods you can use to solve a rational equation. Make a list of the advantages and disadvantages of each method.

Extending

15. Solve $\frac{x^2 - 6x + 5}{x^2 - 2x - 3} = \frac{2 - 3x}{x^2 + 3x + 3}$ correct to two decimal places.
16. Objects A and B move along a straight line. Their positions, s , with respect to an origin, at t seconds, are modelled by the following functions:
- Object A: $s(t) = \frac{7t}{t^2 + 1}$
- Object B: $s(t) = t + \frac{5}{t + 2}$
- When are the objects at the same position?
 - When is object A closer to the origin than object B?

5.5

Solving Rational Inequalities

YOU WILL NEED

- graphing calculator

GOAL

Solve rational inequalities using algebraic and graphical approaches.

rational inequality

a statement that one rational expression is less than or greater than another rational expression
(e.g., $\frac{2x}{x+3} > \frac{x-1}{5x}$)

LEARN ABOUT the Math

The function $P(t) = \frac{20t}{t+1}$ models the population, in thousands, of Nickelford, t years after 1997. The population, in thousands, of nearby New Ironfield is modelled by $Q(t) = \frac{240}{t+8}$.

- ?** How can you determine the time period when the population of New Ironfield exceeded the population of Nickelford?

EXAMPLE 1 Selecting a strategy to solve a problem

Determine the interval(s) of t where the values of $Q(t)$ are greater than the values of $P(t)$.

Solution A: Using an algebraic strategy to solve an inequality

$$\frac{240}{t+8} > \frac{20t}{t+1}$$

The population of New Ironfield exceeds the population of Nickelford when $Q(t) > P(t)$.
 $t \geq 0$ in the context of this problem. There are no other restrictions on the expressions in the rational inequality since the values that make both expressions undefined are negative numbers.

$$\begin{aligned}
 (t+8)(t+1) \left(\frac{240}{t+8} \right) &> (t+8)(t+1) \left(\frac{20t}{t+1} \right) \\
 \cancel{(t+8)} \cancel{(t+1)} \left(\frac{240}{\cancel{t+8}} \right) &> (t+8) \cancel{(t+1)} \left(\frac{20\cancel{t}}{\cancel{t+1}} \right) \\
 240(t+1) &> 20t(t+8) \\
 240t + 240 &> 20t^2 + 160t \\
 0 &> 20t^2 + 160t - 240t - 240 \\
 0 &> 20t^2 - 80t - 240 \\
 0 &> 20(t^2 - 4t - 12) \\
 0 &> 20(t-6)(t+2)
 \end{aligned}$$

Multiply both sides of the inequality by the LCD. The value of the LCD is always positive, since $t \geq 0$, so the inequality sign is unchanged.

Expand and simplify both sides. Then subtract $240t$ and 240 from both sides.

Factor the resulting quadratic expression.

Examine the sign of the factored polynomial expression on the right side of the inequality.

	$t < -2$	$-2 < t < 6$	$t > 6$
$20(t-6)$	-	-	+
$t+2$	-	+	+
$20(t-6)(t+2)$	$(-)(-) = +$	$(-)(+) = -$	$(+)(+) = +$

The inequality $0 > 20(t-6)(t+2)$ is true when the expression on the right side is negative. The sign of the factored quadratic expression changes when $t = -2$ and when $t = 6$, because the expression is zero at these values. Use a table to determine when the sign of the expression is negative on each side of these values.



The inequality $0 > 20(t-6)(t+2)$ is true when $-2 < t < 6$.

The population of New Ironfield exceeded the population of Nickelford for six years after 1997, until 2003.


Since the domain is $t \geq 0$, however, numbers that are negative cannot be included. Therefore, the solution is $0 \leq t < 6$.

Solution B: Solving a rational inequality by graphing two rational functions

To solve $Q(t) > P(t)$, graph $Q(t) = \frac{240}{t+8}$ and $P(t) = \frac{20t}{t+1}$ using graphing technology, and determine the value of t at the intersection point(s).



It helps to bold the graph of $Q(t)$ so you can remember which graph is which. Use window settings that reflect the domain of the functions.



There is only one intersection within the domain of the functions.


From the graphs, $Q(t) > P(t)$ for $0 \leq t < 6$.
The population of New Ironfield exceeded the population of Nickelford until 2003.

If $Q(t) > P(t)$, the graph of $Q(t)$ lies above the graph of $P(t)$.
Looking at the graphs, this is true for the parts of the graph of $Q(t)$ up to the intersection point at $t = 6$.
The graphs will not intersect again because each graph is approaching a different horizontal asymptote.
From the defining equations, the graph of $Q(t)$ is approaching the line $Q = 0$ while the graph of $P(t)$ is approaching the line $P = 20$.

Solution C: Solving a rational inequality by determining the zeros of a combined function

When $Q(t) > P(t)$, $Q(t) - P(t) > 0$.

Graph $f(t) = Q(t) - P(t) = \frac{240}{t+8} - \frac{20t}{t+1}$ and use the zero operation to locate the zero.



Combine the two population functions into a single function, $f(t) = Q(t) - P(t)$. When $Q(t) > P(t)$, $f(t)$ will have positive values.

When a function has positive values, its graph lies above the x-axis.

The graph is above the x-axis for $0 \leq t < 6$.

X	Y1
0	30
2	10.667
4	4
6	0
8	-2.778

By examining the values of $f(t)$ in a table, you can verify that the function continues to decrease but remains positive when $0 \leq t < 6$.

$f(t)$ has positive values for $0 \leq t < 6$.

For the six years after 1997, the population of New Ironfield exceeded the population of Nickelford.

Reflecting

- How is the solution to an inequality different from the solution to an equation?
- In Solution A, how was the rational inequality manipulated to obtain a simpler quadratic inequality?
- In Solution B, how were the graphs of the related rational functions used to find the solution to an inequality?
- In Solution C, how did creating a new function help to solve the inequality?

APPLY the Math

EXAMPLE 2

Selecting a strategy to solve an inequality that involves a linear function and a reciprocal function

Solve $x - 2 < \frac{8}{x}$.

Solution A: Using an algebraic strategy and a sign chart

$$\begin{aligned}
 x - 2 &< \frac{8}{x}, x \neq 0 \\
 x - 2 - \frac{8}{x} &< 0 \\
 \frac{x^2}{x} - \frac{2x}{x} - \frac{8}{x} &< 0 \\
 \frac{x^2 - 2x - 8}{x} &< 0 \\
 \frac{(x - 4)(x + 2)}{x} &< 0
 \end{aligned}$$

Determine any restrictions on x .
Subtract $\frac{8}{x}$ from both sides.

x is the LCD and it can be positive or negative. Multiplying both sides by x would require that two cases be considered, since the inequality sign must be reversed when multiplying by a negative. The alternative is to create an expression with a common denominator, x .

Combine the terms to create a single rational expression.

Factor the numerator.




Examine the sign of the rational expression.

	$x < -2$	$-2 < x < 0$	$0 < x < 4$	$x > 4$
$x - 4$	—	—	—	+
$x + 2$	—	+	+	+
x	—	—	+	+
$\frac{(x - 4)(x + 2)}{x}$	$\frac{(-)(-)}{-} = -$	$\frac{(-)(+)}{-} = +$	$\frac{(-)(+)}{+} = -$	$\frac{(+)(+)}{+} = +$

The sign of a rational expression changes each time the sign of one of its factors changes. Choose a test value in each interval to determine the sign of each part of the expression. Then determine the intervals where the overall expression is negative.

The overall expression is negative when $x < -2$ or when $0 < x < 4$.

The inequality is true when $x \in (-\infty, -2)$ or $x \in (0, 4)$.



Write the solution in interval or set notation, and draw the solution set on a number line.

Solution B: Using graphing technology

$$x - 2 < \frac{8}{x}, x \neq 0$$



Let $f(x) = x - 2$ and $g(x) = \frac{8}{x}$.

The solution set for the inequality will be all x -values for which $f(x) < g(x)$.

Write each side of the inequality as its own function. Enter both functions in the equation editor, using a bold line for $f(x)$.



Graph $f(x)$ and $g(x)$ on the same axes, and use the intersect operation to determine the intersection points.



$f(x) < g(x)$ where the bold graph of $f(x)$ lies beneath the graph of $g(x)$. Notice that the bold linear function is above the reciprocal function on the left side and close to the vertical asymptote, $x = 0$. It is below the reciprocal function on the right side and close to this asymptote.

$f(x) < g(x)$ when $x < -2$ or when $0 < x < 4$.

The solution set is $\{x \in \mathbf{R} \mid x < -2 \text{ or } 0 < x < 4\}$.

You can also use interval notation or a number line to describe the solution set, as in Solution A.

EXAMPLE 3**Determining the solution set for an inequality that involves two rational functions**

Determine the solution set for the inequality $\frac{x+3}{x+1} \geq \frac{x-2}{x-3}$.

Solution A: Using algebra and a sign chart

Rewrite $\frac{x+3}{x+1} \geq \frac{x-2}{x-3}$, $x \neq -1, 3$,

as $\frac{x+3}{x+1} - \frac{x-2}{x-3} \geq 0$.

$$\frac{(x-3)(x+3)}{(x-3)(x+1)} - \frac{(x-2)(x+1)}{(x-3)(x+1)} \geq 0$$

$$\frac{x^2 - 9 - (x^2 - x - 2)}{(x-3)(x+1)} \geq 0$$

$$\frac{x^2 - 9 - x^2 + x + 2}{(x-3)(x+1)} \geq 0$$

$$\frac{x-7}{(x-3)(x+1)} \geq 0$$

Note the restrictions on x .

Subtract $\frac{x-2}{x-3}$ from both sides to create an inequality with zero on the right side.

Subtract the rational expressions on the left side using a common denominator.

Expand and simplify the numerator.

A rational expression is zero when its numerator is zero.

The rational expression is equal to zero when $x = 7$, so 7 is included in the solution set.

Examine the sign of the simplified rational expression on the intervals shown to determine where the rational expression is greater than zero.

	$x < -1$	$-1 < x < 3$	$3 < x < 7$	$x > 7$
$x - 7$	−	−	−	+
$x - 3$	−	−	+	+
$x + 1$	−	+	+	+
$\frac{(x-7)}{(x-3)(x+1)}$	$\frac{-}{(-)(-)} = -$	$\frac{-}{(-)(+)} = +$	$\frac{-}{(+)(+)} = -$	$\frac{+}{(+)(+)} = +$

The expression is undefined at $x = -1$ and $x = 3$. It is equal to 0 at $x = 7$. These numbers create four intervals to consider. Choose a test value in each interval to determine the sign of each part of the expression. Then determine the intervals where the overall expression is positive.

The solution set is $\{x \in \mathbf{R} \mid -1 < x < 3 \text{ or } x \geq 7\}$.



Solution B: Using graphing technology

$$\frac{x+3}{x+1} \geq \frac{x-2}{x-3}, x \neq -1, 3$$




Use each side of the inequality to define a function.
Graph $f(x) = \frac{x+3}{x+1}$ with a bold line and
 $g(x) = \frac{x-2}{x-3}$ with a regular line.

The graph of $f(x)$ has a vertical asymptote at $x = -1$.
The graph for $g(x)$ has a vertical asymptote at $x = 3$.
Both graphs have $y = 1$ as a horizontal asymptote.

Determine the equations of the asymptotes from the equations of the functions.

Use the intersect operation to locate any intersection points.



It looks as though the graphs might intersect on the left side of the screen, as well as on the right side. No matter how far you trace along the left branches, however, you never reach a point where the y -value is the same on both curves.

The functions are equal when $x = 7$.

$f(x) > g(x)$ between the asymptotes at $x = -1$ and $x = 3$, and for $x > 7$.

$f(x) = g(x)$ when $x = 7$.

The solution set for $\frac{x+3}{x+1} \geq \frac{x-2}{x-3}$ is

The bold graph of $f(x)$ is above the graph of $g(x)$ between the two vertical asymptotes and then after the intersection point.



In Summary


Key Ideas

- Solving an inequality means finding all the possible values of the variable that satisfy the inequality.
- To solve a rational inequality algebraically, rearrange the inequality so that one side is zero. Combine the expressions on the no-zero side using a common denominator. Make a table to examine the sign of each factor and the sign of the entire expression on the intervals created by the zeros of the numerator and the denominator.
- Only when you are certain that each denominator is positive can you multiply both sides by the lowest common denominator to make the inequality easier to solve.
- You can always solve a rational inequality using graphing technology.

Need to Know

- When multiplying or dividing both sides of an inequality by a negative it is necessary to reverse the inequality sign to maintain equivalence.
- You can solve an inequality using graphing technology by graphing the functions on each side of the inequality sign and then identifying all the intervals created by the vertical asymptotes and points of intersection. For x -values that satisfy $f(x) > g(x)$, identify the specific intervals where the graph of $f(x)$ is above the graph of $g(x)$. For x -values that satisfy $f(x) < g(x)$, identify the specific intervals where the graph of $f(x)$ is below the graph of $g(x)$.

Consider the following graph:



In this graph, there are four intervals to consider:

$(-\infty, -3)$, $(-3, -2)$, $(-2, 2)$ and $(2, \infty)$. In these intervals, $f(x) > g(x)$ when $x \in (-\infty, -3)$ or $(-2, 2)$, and $f(x) < g(x)$ when $x \in (-3, -2)$ or $(2, \infty)$.

- You can also solve an inequality using graphing technology by creating an equivalent inequality with zero on one side and then identifying the intervals created by the zeros on the graph of the new function. Finding where the graph lies above the x -axis (where $f(x) > 0$) or below the x -axis (where $f(x) < 0$) defines the solutions to the inequality.

CHECK Your Understanding

1. Use the graph shown to determine the solution set for each of the following inequalities.


a) $\frac{x+5}{x-1} < 4$

b) $4x - 1 > \frac{x+5}{x-1}$

2. a) Show that the inequality $\frac{6x}{x+3} \leq 4$ is equivalent to the inequality $\frac{2(x-6)}{(x+3)} \leq 0$.

b) Sketch the solution on a number line.

c) Write the solution using interval notation.



3. a) Show that the inequality $x + 2 > \frac{15}{x}$ is equivalent to the inequality $\frac{(x+5)(x-3)}{x} > 0$.
- b) Use a table to determine the positive/negative intervals for $f(x) = \frac{(x+5)(x-3)}{x}$.
- c) State the solution to the inequality using both set notation and interval notation.

PRACTISING

4. Use algebra to find the solution set for each inequality. Verify your answer using graphing technology.
- a) $\frac{1}{x+5} > 2$ d) $\frac{7}{x-3} \geq \frac{2}{x+4}$
- b) $\frac{1}{2x+10} < \frac{1}{x+3}$ e) $\frac{-6}{x+1} > \frac{1}{x}$
- c) $\frac{3}{x-2} < \frac{4}{x}$ f) $\frac{-5}{x-4} < \frac{3}{x+1}$
5. Use algebra to obtain a factorable expression from each inequality, if necessary. Then use a table to determine interval(s) in which the inequality is true.
- a) $\frac{t^2 - t - 12}{t-1} < 0$ d) $t-1 < \frac{30}{5t}$
- b) $\frac{t^2 + t - 6}{t-4} \geq 0$ e) $\frac{2t-10}{t} > t+5$
- c) $\frac{6t^2 - 5t + 1}{2t+1} > 0$ f) $\frac{-t}{4t-1} \geq \frac{2}{t-9}$
6. Use graphing technology to solve each inequality.
- a) $\frac{x+3}{x-4} \geq \frac{x-1}{x+6}$ d) $\frac{x}{x+9} \geq \frac{1}{x+1}$
- b) $x+5 < \frac{x}{2x+6}$ e) $\frac{x-8}{x} > 3-x$
- c) $\frac{x}{x+4} \leq \frac{1}{x+1}$ f) $\frac{x^2-16}{(x-1)^2} \geq 0$
7. a) Find all the values of x that make the following inequality true:
K $\frac{3x-8}{2x-1} > \frac{x-4}{x+1}$
- b) Graph the solution set on a number line. Write the solution set using interval notation and set notation.

8. a) Use an algebraic strategy to solve the inequality $\frac{-6t}{t-2} < \frac{-30}{t-2}$.
 b) Graph both inequalities to verify your solution.
 c) Can these rational expressions be used to model a real-world situation? Explain.
9. The equation $f(t) = \frac{5t}{t^2 + 3t + 2}$ models the bacteria count, in thousands, for a sample of tap water that is left to sit over time, t , in days. The equation $g(t) = \frac{15t}{t^2 + 9}$ models the bacteria count, in thousands, for a sample of pond water that is also left to sit over several days. In both models, $t > 0$. Will the bacteria count for the tap water sample ever exceed the bacteria count for the pond water? Justify your answer.
10. Consider the inequality $0.5x - 2 < \frac{5}{2x}$.
 a) Rewrite the inequality so that there is a single, simplified expression on one side and a zero on the other side.
 b) List all the factors of the rational expression in a table, and determine on which intervals the inequality is true.
11. An economist for a sporting goods company estimates the revenue and cost functions for the production of a new snowboard. These functions are $R(x) = -x^2 + 10x$ and $C(x) = 4x + 5$, respectively, where x is the number of snowboards produced, in thousands. The average profit is defined by the function $AP(x) = \frac{P(x)}{x}$, where $P(x)$ is the profit function. Determine the production levels that make $AP(x) > 0$.
12. a) Explain why the inequalities $\frac{x+1}{x-1} < \frac{x+3}{x+2}$ and $\frac{x+5}{(x-1)(x+2)} < 0$ are equivalent.
 b) Describe how you would use a graphing calculator to solve these inequalities.
 c) Explain how you would use a table to solve these inequalities.

Extending

13. Solve $|\frac{x}{x-4}| \geq 1$.
14. Solve $\frac{1}{\sin x} < 4$, $0^\circ \leq x \leq 360^\circ$.
15. Solve $\frac{\cos(x)}{x} > 0.5$, $0^\circ < x < 90^\circ$.